

t11_jordan18 (TMZiA- DeM5fG8GZ7jDNmyXsujp6YVeDswjKN)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k1_jordan2c : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_topreal1 : \iota \Rightarrow \iota$ be given. Let $k2_jordan2c : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xboole_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_topreal1 : \iota \Rightarrow \iota$ be given. Let $k5_topreal1 : \iota \Rightarrow \iota$ be given. Let $k6_topreal1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_xxreal_0 : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (\neg(\neg r1_xboole_0 X0 X1) \wedge (\forall X2. \neg(X2 \in X0) \wedge (X2 \in X1))) \wedge (\neg(\exists X2. (X2 \in X0) \wedge (X2 \in X1))) \wedge (r1_xboole_0 X0 X1) \quad (1)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow ((X0 \in k7_topreal1 X0) \wedge ((X0 \in k5_topreal1 X0) \wedge ((X0 \in k4_topreal1 X0) \wedge (X0 \in k6_topreal1 X0)))) \quad (2)$$

Assume the following.

$$\forall X0. (m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 (k15_euclid X0)))) \Rightarrow (r1_xboole_0 (k1_jordan2c X0 X1) (k2_jordan2c X0 X1))) \quad (3)$$

Assume the following.

$$((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (5)$$

Assume the following.

$$k1_xxreal_0 = k1_numbers \quad (6)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (\neg(X1 \in k1_jordan2c np_2 X0) \wedge (r1_tarski (k4_topreal1 X1) (k2_jordan2c np_2 X0)))$$