

t11_jordan5d

(TMHoP1rKQPwyiFzMFYBjvAXkhuzbocmwbpa)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_6 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_topreal1 : \iota \Rightarrow o$ be given. Let $v2_topreal1 : \iota \Rightarrow o$ be given. Let $v1_goboard5 : \iota \Rightarrow o$ be given. Let $v2_goboard5 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k9_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k3_topreal1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $np_4 : \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_compts_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k8_pscomp_1 : \iota \Rightarrow \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \neg (X0 \in X1) \wedge (v1_xboole_0 X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (2)$$

Assume the following.

$$\forall X0. ((\neg v3_funct_1 X0) \wedge ((\neg v1_xboole_0 X0) \wedge ((v1_finseq_6 X0 (u1_struct_0 (k15_euclid np_2))) \wedge ((v1_topreal1 X0) \wedge ((v2_topreal1 X0) \wedge ((v1_goboard5 X0) \wedge ((v2_goboard5 X0) \wedge (m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2)))))))))) \Rightarrow (\neg r1_xxreal_0 (k3_finseq_1 X0) np_4) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ & (\forall X1.(v7_ordinal_1 X1) \Rightarrow ((X1 \in k1_relset_1 k5_numbers X0) \Leftrightarrow \\ & ((r1_xxreal_0 np_1 X1) \wedge (r1_xxreal_0 X1 (k3_finseq_1 X0)))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.((\neg v1_xboole_0 X1) \wedge ((v2_compts_1 X1 (k15_euclid \\ & np_2)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ & np_2)))))) \Rightarrow ((X0 \in X1) \Rightarrow ((r1_xxreal_0 (k6_pscomp_1 X1) (k17_euclid \\ & X0)) \wedge ((r1_xxreal_0 (k17_euclid X0) (k8_pscomp_1 X1)) \wedge ((r1_xxreal_0 \\ & (k9_pscomp_1 X1) (k18_euclid X0)) \wedge (r1_xxreal_0 (k18_euclid X0) \\ & (k7_pscomp_1 X1)))))) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (((X1 \in k4_finseq_1 \\ & X0) \wedge (r1_xxreal_0 np_2 (k3_finseq_1 X0))) \Rightarrow (k7_partfun1 (u1_struct_0 \\ & (k15_euclid np_2)) X0 X1 \in k3_topreal1 np_2 X0))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_4) \wedge (m2_subset_1 np_4 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_4 k5_numbers) \wedge (m1_subset_1 np_4 k1_numbers)) \end{aligned} \quad (9)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (10)$$

Assume the following.

$$r1_xxreal_0 np_2 np_4 \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Leftrightarrow(m1_finseq_1 X1 X0) \quad (13)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (14)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (16)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (17)$$

Assume the following.

$$\forall X0.(m1_finseq_1 X0 (u1_struct_0 (k15_euclid np_2)))\Rightarrow (v2_compts_1 (k3_topreal1 np_2 X0) (k15_euclid np_2)) \quad (18)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (20)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (21)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (22)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(m1_finseq_1 X1 (u1_struct_0 (k15_euclid X0))))\Rightarrow(m1_subset_1 (k3_topreal1 X0 X1) (k1_zfmisc_1 (u1_struct_0 (k15_euclid X0)))) \quad (23)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow (m2_subset_1 (k3_finseq_1 X0) k1_numbers k5_numbers) \quad (24)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0)\wedge(v1_xxreal_0 X1))\Rightarrow (r1_xxreal_0 X0 X1)\vee(r1_xxreal_0 X1 X0) \quad (25)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (26)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow ((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))) \quad (27)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (28)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (29)$$

Theorem 1

$$\forall X0.((\neg v1_xboole_0 X0)\wedge((\neg v3_funct_1 X0)\wedge((v1_finseq_6 X0 (u1_struct_0 (k15_euclid np_2)))\wedge((v1_topreal1 X0)\wedge((v2_topreal1 X0)\wedge((v1_goboard5 X0)\wedge((v2_goboard5 X0)\wedge(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))))))))))\Rightarrow(\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers)\Rightarrow(((r1_xxreal_0 np_1 X1)\wedge(r1_xxreal_0 X1 (k3_finseq_1 X0))\Rightarrow((r1_xxreal_0 (k9_pscomp_1 (k3_topreal1 np_2 X0)) (k18_euclid (k7_partfun1 (u1_struct_0 (k15_euclid np_2)) X0 X1)))\wedge(r1_xxreal_0 (k18_euclid (k7_partfun1 (u1_struct_0 (k15_euclid np_2)) X0 X1)) (k7_pscomp_1 (k3_topreal1 np_2 X0)))))))$$