

# t11\_matrlin (TMWQfTEJSfU- UGFc3TRxoBeq1uZRgbhCunbA)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l2\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k7\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k4\_rlvect\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k4\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (1)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (2)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k4\_finseq\_1 X0 = k9\_xtuple\_0 X0) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v3\_rlvect\_1 \\ & X0) \wedge ((v4\_rlvect\_1 X0) \wedge (l2\_algstr\_0 X0)))) \Rightarrow (\forall X1. (m2\_finseq\_1 \\ & X1 (u1\_struct\_0 X0)) \Rightarrow ((\forall X2. (m1\_subset\_1 X2 k5\_numbers) \Rightarrow \\ & ((X2 \in k4\_finseq\_1 X1) \Rightarrow (k1\_funct\_1 X1 X2 = k4\_struct\_0 X0))) \Rightarrow (k4\_rlvect\_1 \\ & X0 X1 = k4\_struct\_0 X0))) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(m2\_finseq\_1 X1 X0)\Rightarrow((v1\_funct\_1 X1)\wedge((v1\_finseq\_1 X1)\wedge(m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers X0)))))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(m1\_finseq\_1 X1 X0)\Rightarrow((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge(v1\_finseq\_1 X1))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge((v5\_relat\_1 X1 X0)\wedge(v1\_funct\_1 X1)))\Rightarrow(\forall X2.(X2 \in k9\_xtuple\_0 X1)\Rightarrow(k7\_partfun1 X0 X1 X2 = k1\_funct\_1 X1 X2)) \quad (7)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (9)$$

**Theorem 1**

$$\forall X0.((\neg v2\_struct\_0 X0)\wedge((v13\_algstr\_0 X0)\wedge((v3\_rlvect\_1 X0)\wedge((v4\_rlvect\_1 X0)\wedge(l2\_algstr\_0 X0)))))\Rightarrow(\forall X1.(m2\_finseq\_1 X1 (u1\_struct\_0 X0))\Rightarrow((\forall X2.(v7\_ordinal1 X2)\Rightarrow((X2 \in k9\_xtuple\_0 X1)\Rightarrow(k7\_partfun1 (u1\_struct\_0 X0) X1 X2 = k4\_struct\_0 X0)))\Rightarrow(k4\_rlvect\_1 X0 X1 = k4\_struct\_0 X0)))$$