

t11_mesfun6c
(TMTQYyabE4foaPeAMkQ371yvw3Kk64cccKK)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_mesfun6c : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_comseq_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_comseq_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_mesfunc6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((v1_funct_1 X1) \wedge (\\ & \quad m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \Rightarrow (\\ & \quad \forall X2. ((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 \\ & \quad X0 k2_numbers)))) \Rightarrow ((r2_relset_1 X0 k1_numbers (k5_comseq_3 X0 \\ & \quad (k2_valued_1 X0 k2_numbers k2_numbers X1 X2)) (k3_valued_1 X0 k1_numbers \\ & \quad k1_numbers (k5_comseq_3 X0 X1) (k5_comseq_3 X0 X2))) \wedge (r2_relset_1 \\ & \quad X0 k1_numbers (k6_comseq_3 X0 (k2_valued_1 X0 k2_numbers k2_numbers \\ & \quad X1 X2)) (k3_valued_1 X0 k1_numbers k1_numbers (k6_comseq_3 X0 X1) \\ & \quad (k6_comseq_3 X0 X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. ((\neg v1_xboole_0 X1) \wedge \\ & \quad ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\ & \quad (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2. ((v1_funct_1 X2) \wedge (m1_subset_1 \\ & \quad X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \Rightarrow (\forall X3. (\\ & \quad (v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 \\ & \quad k1_numbers)))) \Rightarrow (\forall X4. (m2_subset_1 X4 (k1_zfmisc_1 X0) \\ & \quad X1) \Rightarrow (((r1_mesfunc6 X0 X1 X2 X4) \wedge (r1_mesfunc6 X0 X1 X3 X4)) \Rightarrow (r1_mesfunc6 \\ & \quad X0 X1 (k3_valued_1 X0 k1_numbers k1_numbers X2 X3) X4)))))) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((r2_relset_1 X0 X1 X2 X3)\Leftrightarrow(X2 = X3)) \quad (3)$$

Assume the following.

$$v3_membered k1_numbers \quad (4)$$

Assume the following.

$$v1_membered k2_numbers \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers))))\Rightarrow((v1_funct_1 (k6_comseq_3 X0 X1))\wedge(m1_subset_1 (k6_comseq_3 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers))))\Rightarrow((v1_funct_1 (k5_comseq_3 X0 X1))\wedge(m1_subset_1 (k5_comseq_3 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.((v3_membered X1)\wedge((v3_membered X2)\wedge(((v1_funct_1 X3)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\wedge((v1_funct_1 X4)\wedge(m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 X2)))))))\Rightarrow((v1_funct_1 (k3_valued_1 X0 X1 X2 X3 X4))\wedge(m1_subset_1 (k3_valued_1 X0 X1 X2 X3 X4) (k1_zfmisc_1 (k2_zfmisc_1 X0 k1_numbers)))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.\forall X4.((v1_membered X1)\wedge((v1_membered X2)\wedge(((v1_funct_1 X3)\wedge(m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\wedge((v1_funct_1 X4)\wedge(m1_subset_1 X4 (k1_zfmisc_1 (k2_zfmisc_1 X0 X2)))))))\Rightarrow((v1_funct_1 (k2_valued_1 X0 X1 X2 X3 X4))\wedge(m1_subset_1 (k2_valued_1 X0 X1 X2 X3 X4) (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \quad (9)$$

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge (m1_subset_1 \\
& X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \Rightarrow (\forall X3.(\\
& m2_subset_1 X3 (k1_zfmisc_1 X0) X1) \Rightarrow ((r1_mesfun6c X0 X1 X2 X3) \Leftrightarrow \\
& ((r1_mesfunc6 X0 X1 (k5_comseq_3 X0 X2) X3) \wedge (r1_mesfunc6 X0 X1 (\\
& k6_comseq_3 X0 X2) X3))))))
\end{aligned} \tag{10}$$

Theorem 1

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((\neg v1_xboole_0 X1) \wedge \\
& ((v1_prob_1 X1 X0) \wedge ((v4_prob_1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 \\
& (k1_zfmisc_1 X0)))))) \Rightarrow (\forall X2.((v1_funct_1 X2) \wedge (m1_subset_1 \\
& X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \Rightarrow (\forall X3.(\\
& (v1_funct_1 X3) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 \\
& k2_numbers)))) \Rightarrow (\forall X4.(m2_subset_1 X4 (k1_zfmisc_1 X0) \\
& X1) \Rightarrow (((r1_mesfun6c X0 X1 X2 X4) \wedge (r1_mesfun6c X0 X1 X3 X4)) \Rightarrow (r1_mesfun6c \\
& X0 X1 (k2_valued_1 X0 k2_numbers k2_numbers X2 X3) X4))))))
\end{aligned}$$