

t11_monoid_0 (TM-
RCtNK6PcjPafUyRmxWDW8rztH5BLocDPJ)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l3_algstr_0 : \iota \Rightarrow o$ be given. Let $v14_monoid_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k6_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $u2_algstr_0 : \iota \Rightarrow \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v6_monoid_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_binop_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\neg v1_xboole_0 (u1_struct_0 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(l3_algstr_0 X0) \Rightarrow ((v1_funct_1 (u2_algstr_0 X0)) \wedge ((v1_funct_2 (u2_algstr_0 X0) (k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 X0)) (u1_struct_0 X0)) \wedge (m1_subset_1 (u2_algstr_0 X0) (k1_zfmisc_1 (k2_zfmisc_1 (k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 X0)) (u1_struct_0 X0)))))) \quad (2)$$

Assume the following.

$$\forall X0.(l3_algstr_0 X0) \Rightarrow (l1_struct_0 X0) \quad (3)$$

Assume the following.

$$\forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.((v1_funct_1 X1) \wedge (v1_funct_2 X1 (k2_zfmisc_1 X0 X0) X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 (k2_zfmisc_1 X0 X0) X0)))) \Rightarrow ((v6_monoid_0 X1 X0) \Leftrightarrow (\forall X2.(m1_subset_1 X2 X0) \Rightarrow (\forall X3.(m1_subset_1 X3 X0) \Rightarrow (\forall X4.(m1_subset_1 X4 X0) \Rightarrow ((k5_binop_1 X0 X1 X2 X3 = k5_binop_1 X0 X1 X2 X4) \Rightarrow (X3 = X4)))))))) \quad (4)$$

Assume the following.

$$\forall X0.(l3_algstr_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (k6_algstr_0 X0 X1 X2 = k5_binop_1 (u1_struct_0 X0) (u2_algstr_0 X0) X1 X2))) \quad (5)$$

Assume the following.

$$\forall X0.((\neg v2_struct_0 X0) \wedge (l3_algstr_0 X0)) \Rightarrow ((v14_monoid_0 X0) \Leftrightarrow (v6_monoid_0 (u2_algstr_0 X0) (u1_struct_0 X0))) \quad (6)$$

Theorem 1

$$\begin{aligned} \forall X0.((\neg v2_struct_0 X0) \wedge (l3_algstr_0 X0)) \Rightarrow ((v14_monoid_0 X0) \Leftrightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 \\ (u1_struct_0 X0)) \Rightarrow ((k6_algstr_0 X0 X1 X2 = k6_algstr_0 X0 X1 X3) \Rightarrow \\ (X2 = X3))))))) \end{aligned}$$