

t11_nfcont_3

(TMS1H1UMpaJEnC9qyFGDLSvfXSxwPCvzjyX)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_normsp_0 : \iota \Rightarrow o$ be given. Let $v4_normsp_0 : \iota \Rightarrow o$ be given. Let $v2_normsp_1 : \iota \Rightarrow o$ be given. Let $l1_normsp_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m1_rcomp_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $r1_nfcont_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k9_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_nfcont_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow ((X0 \in k9_xtuple_0 X1) \Rightarrow (k9_relat_1 X1 X0 = k1_tarski (k1_funct_1 X1 X0))) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge ((v3_normsp_0 X0) \wedge ((v4_normsp_0 X0) \wedge ((v2_normsp_1 X0) \wedge (l1_normsp_1 X0)))))))))) \Rightarrow \\ & (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. (m1_nfcont_1 X2 X0 X1) \Rightarrow (X1 \in X2))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski (k1_tarski X0) X1) \Leftrightarrow (X0 \in X1) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X1)\Rightarrow(k7_relat_1 X1 X0 = k7_relat_1 X1 (k3_xboole_0 (k9_xtuple_0 X1) X0)) \quad (4)$$

Assume the following.

$$\begin{aligned} &\forall X0.((\neg v2_struct_0 X0)\wedge((v13_algstr_0 X0)\wedge((v2_rlvect_1 X0)\wedge((v3_rlvect_1 X0)\wedge((v4_rlvect_1 X0)\wedge((v5_rlvect_1 X0)\wedge \\ &((v6_rlvect_1 X0)\wedge((v7_rlvect_1 X0)\wedge((v8_rlvect_1 X0)\wedge((v3_normsp_0 X0)\wedge((v4_normsp_0 X0)\wedge((v2_normsp_1 X0)\wedge(l1_normsp_1 X0))))))))))\Rightarrow \\ &(\forall X1.((v1_funct_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers (u1_struct_0 X0))))\Rightarrow(\forall X2.(v1_xreal_0 X2)\Rightarrow \\ &((r1_nfcont_3 X0 X1 X2)\Leftrightarrow((X2 \in k1_relset_1 k1_numbers X1)\wedge(\forall X3. (m1_nfcont_1 X3 X0 (k7_partfun1 (u1_struct_0 X0) X1 X2))\Rightarrow(\exists X4. (m1_rcomp_1 X4 X2)\wedge(r1_tarski (k7_relset_1 k1_numbers (u1_struct_0 X0) X1 X4) X3))))))))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.r1_tarski X0 X0 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 X0))\Rightarrow(k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(k7_relset_1 X0 X1 X2 X3 = k7_relat_1 X2 X3) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.(m1_rcomp_1 X1 X0)\Rightarrow(m1_subset_1 X1 (k1_zfmisc_1 k1_numbers))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_relat_1 X1)\wedge((v5_relat_1 X1 X0)\wedge(v1_funct_1 X1)))\Rightarrow(m1_subset_1 (k7_partfun1 X0 X1 X2) X0) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge((v5_relat_1 X1 X0)\wedge(v1_funct_1 X1)))\Rightarrow(\forall X2.(X2 \in k9_xtuple_0 X1)\Rightarrow(k7_partfun1 X0 X1 X2 = k1_funct_1 X1 X2)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k3_xboole_0 X0 X1)\Leftrightarrow(\forall X3.(X3 \in X2)\Leftrightarrow((X3 \in X0)\wedge(X3 \in X1))) \quad (13)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0)\Rightarrow(\forall X1.k9_relat_1 X0 X1 = k7_relat_1 X0 (k1_tarski X1)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (16)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(\forall X1.((\neg v2_struct_0 X1)\wedge(v13_algstr_0 X1)\wedge((v2_rlvect_1 X1)\wedge((v3_rlvect_1 X1)\wedge((v4_rlvect_1 X1)\wedge((v5_rlvect_1 X1)\wedge((v6_rlvect_1 X1)\wedge((v7_rlvect_1 X1)\wedge((v8_rlvect_1 X1)\wedge((v3_normsp_0 X1)\wedge((v4_normsp_0 X1)\wedge((v2_normsp_1 X1)\wedge(l1_normsp_1 X1))))))))))))))\Rightarrow(\forall X2.((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers (u1_struct_0 X1))))))\Rightarrow((\exists X3.(m1_rcomp_1 X3 X0)\wedge(k9_subset_1 k1_numbers (k1_relset_1 k1_numbers X2) X3 = k1_tarski X0))\Rightarrow(r1_nfcont_3 X1 X2 X0))))$$