

# t11\_qc\_lang2 (TMTE- HDz8QuxkH9kfP5HcDN3J8885ndP8wxC)

October 27, 2020

Let  $m1\_qc\_lang1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_qc\_lang1 : \iota \Rightarrow \iota$  be given. Let  $k2\_qc\_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k19\_qc\_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k14\_qc\_lang1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k20\_qc\_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k18\_qc\_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k13\_qc\_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k9\_qc\_lang1 \\ X0)) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (k9\_qc\_lang1 X0)) \Rightarrow ((k19\_qc\_lang1 \\ X0 (k14\_qc\_lang1 X0 X1 X2) = X1) \wedge (k20\_qc\_lang1 X0 (k14\_qc\_lang1 \\ X0 X1 X2) = X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k9\_qc\_lang1 \\ X0)) \Rightarrow (k18\_qc\_lang1 X0 (k13\_qc\_lang1 X0 X1) = X1)) \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1\_qc\_lang1 X0) \wedge ((m1\_subset\_1 \\ X1 (k9\_qc\_lang1 X0)) \wedge (m1\_subset\_1 X2 (k9\_qc\_lang1 X0)))) \Rightarrow (m1\_subset\_1 \\ (k14\_qc\_lang1 X0 X1 X2) (k9\_qc\_lang1 X0)) \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.((m1\_qc\_lang1 X0) \wedge (m1\_subset\_1 X1 (k9\_qc\_lang1 \\ X0))) \Rightarrow (m1\_subset\_1 (k13\_qc\_lang1 X0 X1) (k9\_qc\_lang1 X0)) \tag{4}$$

Assume the following.

$$\forall X0.(m1\_qc\_lang1 X0) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k9\_qc\_lang1 \\ X0)) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (k9\_qc\_lang1 X0)) \Rightarrow (k2\_qc\_lang2 \\ X0 X1 X2 = k13\_qc\_lang1 X0 (k14\_qc\_lang1 X0 X1 (k13\_qc\_lang1 X0 X2)))))) \tag{5}$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(m1\_qc\_lang1\ X0) \Rightarrow (\forall X1.(m1\_subset\_1\ X1\ (k9\_qc\_lang1 \\ & \quad X0)) \Rightarrow (\forall X2.(m1\_subset\_1\ X2\ (k9\_qc\_lang1\ X0)) \Rightarrow (\forall X3. \\ & \quad (m1\_subset\_1\ X3\ (k9\_qc\_lang1\ X0)) \Rightarrow (\forall X4.(m1\_subset\_1\ X4 \\ & \quad (k9\_qc\_lang1\ X0)) \Rightarrow ((k2\_qc\_lang2\ X0\ X1\ X2 = k2\_qc\_lang2\ X0\ X3\ X4) \Rightarrow \\ & \quad ((X1 = X3) \wedge (X2 = X4)))))) \end{aligned}$$