

# t11\_relset\_1 (TMZr- cQQ8q9Rh9RHoADRAMQD9Nm5unKsKgRH)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k2\_relat\_1 : \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_relat\_1 X0) \Rightarrow ((k10\_xtuple\_0 X0 = k9\_xtuple\_0 (k2\_relat\_1 X0)) \wedge (k9\_xtuple\_0 X0 = k10\_xtuple\_0 (k2\_relat\_1 X0))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (k3\_relset\_1 X0 X1 X2 = k2\_relat\_1 X2) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1) \wedge (v5\_relat\_1 X1 X0)) \Rightarrow (k2\_relset\_1 X0 X1 = k10\_xtuple\_0 X1) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (m1\_subset\_1 (k3\_relset\_1 X0 X1 X2) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 X0))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v4\_relat\_1 X2 X0) \wedge (v5\_relat\_1 X2 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (7)$$

**Theorem 1**

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((k1\_relset\_1 X1 (k3\_relset\_1 X0 X1 X2) = k2\_relset\_1 X1 X2)\wedge(k2\_relset\_1 X0 (k3\_relset\_1 X0 X1 X2) = k1\_relset\_1 X0 X2))$$