

t11\_scmring1  
(TMaSRG1yYXSguJJtRQeJ6gkxVgThoHV2SvA)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $l1\_struct\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_card\_3 : \iota \Rightarrow \iota$  be given. Let  $k3\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_ami\_2 : \iota$  be given. Let  $k1\_scmring1 : \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_ami\_2 : \iota$  be given. Let  $k2\_ami\_2 : \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_scmring1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k16\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k7\_funcop\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v4\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_2 : \iota$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. k1\_funct\_1 (k16\_funcop\_1 X0 X1) X0 = X1 \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (\forall X2. \\ & ((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow ((X0 \in k9\_xtuple\_0 X1) \Rightarrow (k1\_funct\_1 \\ & (k1\_funct\_4 X2 X1) X0 = k1\_funct\_1 X1 X0))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (k9\_xtuple\_0 (k2\_funcop\_1 X0 X1) = X0) \wedge (r1\_tarski (k10\_xtuple\_0 (k2\_funcop\_1 X0 X1)) (k1\_tarski X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k7\_funcop\_1 X0 X1 = k2\_funcop\_1 X0 X1 \quad (4)$$

Assume the following.

$$\forall X0. ((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (v4\_funct\_1 (k4\_card\_3 X0)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(((v1\_relat\_1 X0)\wedge(v1\_funct\_1 X0))\wedge((v1\_relat\_1 X1)\wedge(v1\_funct\_1 X1)))\Rightarrow((v1\_relat\_1 (k3\_relat\_1 X0 X1))\wedge(v1\_funct\_1 (k3\_relat\_1 X0 X1))) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v1\_relat\_1 (k2\_funcop\_1 X0 X1))\wedge(v1\_funct\_1 (k2\_funcop\_1 X0 X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.v1\_relat\_1 (k3\_relat\_1 X0 X1) \quad (8)$$

Assume the following.

$$(v1\_funct\_1 k3\_ami\_2)\wedge((v1\_funct\_2 k3\_ami\_2 k1\_ami\_2 np\_2)\wedge(m1\_subset\_1 k3\_ami\_2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k1\_ami\_2 np\_2)))) \quad (9)$$

Assume the following.

$$\forall X0.((\neg v2\_struct\_0 X0)\wedge(l1\_struct\_0 X0))\Rightarrow((v1\_relat\_1 (k1\_scmring1 X0))\wedge((v4\_relat\_1 (k1\_scmring1 X0) np\_2)\wedge((v1\_funct\_1 (k1\_scmring1 X0))\wedge(v1\_partfun1 (k1\_scmring1 X0) np\_2)))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.k16\_funcop\_1 X0 X1 = k7\_funcop\_1 (k1\_tarski X0) X1 \quad (11)$$

Assume the following.

$$\forall X0.((\neg v2\_struct\_0 X0)\wedge(l1\_struct\_0 X0))\Rightarrow(\forall X1.(m1\_subset\_1 X1 (k4\_card\_3 (k3\_relat\_1 k3\_ami\_2 (k1\_scmring1 X0))))\Rightarrow(\forall X2.(m2\_subset\_1 X2 k1\_ami\_2 k2\_ami\_2)\Rightarrow(\forall X3.(m1\_subset\_1 X3 (u1\_struct\_0 X0))\Rightarrow(k4\_scmring1 X0 X1 X2 X3 = k1\_funct\_4 X1 (k16\_funcop\_1 X2 X3)))))) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k1\_tarski X0)\Leftrightarrow(\forall X2.(X2 \in X1)\Leftrightarrow(X2 = X0)) \quad (13)$$

Assume the following.

$$\forall X0.(v4\_funct\_1 X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 X0)\Rightarrow((v1\_relat\_1 X1)\wedge(v1\_funct\_1 X1))) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (15)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge (l1\_struct\_0 X0)) \Rightarrow (\forall X1. \\ & (m1\_subset\_1 X1 (k4\_card\_3 (k3\_relat\_1 k3\_ami\_2 (k1\_scmring1 \\ & X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 X2 k1\_ami\_2 k2\_ami\_2) \Rightarrow (\forall X3. \\ & (m1\_subset\_1 X3 (u1\_struct\_0 X0)) \Rightarrow (k1\_funct\_1 (k4\_scmring1 X0 \\ & X1 X2 X3) X2 = X3)))) \end{aligned}$$