

t11_seqm_3
(TMJK9mMwSyL8CB3Hog4qpUP29WnNSd6Ftfo)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_valued_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k5_complex1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_xxreal_0 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0. ((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k1_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers)))))) \Rightarrow \\ & ((v7_valued_0 X0) \Leftrightarrow (\forall X1. (m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow \\ & (\forall X2. (m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow ((r1_xxreal_0 \\ & X1 X2) \Rightarrow (r1_xxreal_0 (k8_nat_1 k1_numbers X0 X1) (k8_nat_1 k1_numbers \\ & X0 X2))))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (r1_xxreal_0 k6_numbers X0) \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \tag{3}$$

Assume the following.

$$k6_numbers = k1_xboole_0 \tag{4}$$

Assume the following.

$$k5_numbers = k4_ordinal1 \tag{5}$$

Assume the following.

$$k5_complex1 = k1_xboole_0 \quad (6)$$

Assume the following.

$$(\neg v1_xboole_0 \ k4_ordinal1) \wedge (v3_ordinal1 \ k4_ordinal1) \quad (7)$$

Assume the following.

$$m2_subset_1 \ k6_numbers \ k1_numbers \ k5_numbers \quad (8)$$

Assume the following.

$$m1_subset_1 \ k5_numbers \ (k1_zfmisc_1 \ k1_numbers) \quad (9)$$

Assume the following.

$$k1_xxreal_0 = k1_numbers \quad (10)$$

Assume the following.

$$\forall X0. (m1_subset_1 \ X0 \ k4_ordinal1) \Rightarrow (v7_ordinal1 \ X0) \quad (11)$$

Assume the following.

$$\begin{aligned} \forall X0. (v1_xboole_0 \ X0) \Rightarrow (\forall X1. (m1_subset_1 \ X1 \ (k1_zfmisc_1 \\ X0)) \Rightarrow (v1_xboole_0 \ X1)) \end{aligned} \quad (12)$$

Theorem 1

$$\begin{aligned} \forall X0. ((v1_funct_1 \ X0) \wedge ((v1_funct_2 \ X0 \ k5_numbers \ k1_numbers) \wedge \\ (m1_subset_1 \ X0 \ (k1_zfmisc_1 \ (k2_zfmisc_1 \ k5_numbers \ k1_numbers)))))) \Rightarrow \\ ((v7_valued_0 \ X0) \Rightarrow (\forall X1. (m2_subset_1 \ X1 \ k1_numbers \ k5_numbers) \Rightarrow \\ (r1_xxreal_0 \ (k8_nat_1 \ k1_numbers \ X0 \ k6_numbers) \ (k8_nat_1 \ k1_numbers \\ X0 \ X1)))) \end{aligned}$$