

## t11\_sin\_cos7

(TMJQCwWkM7ANotsVoQwhg6Pb4EWaNRjFaHd)

October 27, 2020

Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $k3\_square\_1 : \iota \Rightarrow \iota$  be given. Let  $k9\_binop\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k10\_binop\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xcmplx\_0 : \iota \Rightarrow \iota$  be given. Let  $k6\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow ((\neg r1\_xxreal\_0 np\_1 (k3\_square\_1 X0)) \Rightarrow ((\neg r1\_xxreal\_0 X0 (k4\_xcmplx\_0 np\_1)) \wedge (\neg r1\_xxreal\_0 np\_1 X0))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\neg(\neg r1\_xxreal\_0 np\_1 X0) \wedge (r1\_xxreal\_0 (k6\_xcmplx\_0 np\_1 X0) k6\_numbers)) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (\neg(\neg r1\_xxreal\_0 X0 (k4\_xcmplx\_0 np\_1)) \wedge (r1\_xxreal\_0 (k2\_xcmplx\_0 np\_1 X0) k6\_numbers)) \quad (3)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X1)) \Rightarrow (k9\_binop\_2 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0) \wedge (v1\_xreal\_0 X1)) \Rightarrow (k10\_binop\_2 X0 X1 = k6\_xcmplx\_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(k9\_binop\_2 X0 X1 = k9\_binop\_2 X1 X0) \quad (7)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xreal\_0 X0) \quad (8)$$

**Theorem 1**

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow((\neg r1\_xxreal\_0 np\_1 (k3\_square\_1 X0))\Rightarrow((\neg r1\_xxreal\_0 (k9\_binop\_2 X0 np\_1) k6\_numbers)\wedge(\neg r1\_xxreal\_0 (k10\_binop\_2 np\_1 X0) k6\_numbers)))$$