

t11_tdlat_2 (TMY-
eEjN6StGBDT3NqsdjrWUnVfW8Mz1UCGm)

October 27, 2020

Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_pcomps_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k7_subset_1 X0 X1 X2 = k4_xboole_0 X1 X2) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (m1_subset_1 (k7_subset_1 X0 X1 X2) (k1_zfmisc_1 X0)) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((l1_pre_topc X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 (u1_struct_0 X0)))))) \Rightarrow (m1_subset_1 (k1_pcomps_1 X0 X1) (k1_zfmisc_1 (k1_zfmisc_1 (u1_struct_0 X0)))) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k4_xboole_0 X0 X1) \Leftrightarrow (\forall X3.(X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (\neg X3 \in X1))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski X0 X1) \Leftrightarrow (\forall X2.(X2 \in X0) \Rightarrow (X2 \in X1)) \quad (6)$$

Assume the following.

$$\begin{aligned}
& \forall X0.(l1_pre_topc\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1 \\
& \quad (k1_zfmisc_1\ (u1_struct_0\ X0)))) \Rightarrow (\forall X2.(m1_subset_1\ X2 \\
& \quad (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0)))) \Rightarrow ((X2 = k1_pcomps_1 \\
& \quad X0\ X1) \Leftrightarrow (\forall X3.(m1_subset_1\ X3\ (k1_zfmisc_1\ (u1_struct_0 \\
& \quad X0))) \Rightarrow ((X3 \in X2) \Leftrightarrow (\exists X4.(m1_subset_1\ X4\ (k1_zfmisc_1\ (u1_struct_0 \\
& \quad X0))) \wedge ((X3 = k2_pre_topc\ X0\ X4) \wedge (X4 \in X1))))))
\end{aligned} \tag{7}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((v2_pre_topc\ X0) \wedge (l1_pre_topc\ X0)) \Rightarrow (\forall X1. \\
& \quad (m1_subset_1\ X1\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0\ X0)))) \Rightarrow \\
& \quad (\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k1_zfmisc_1\ (u1_struct_0 \\
& \quad X0)))) \Rightarrow (r1_tarSKI\ (k7_subset_1\ (k1_zfmisc_1\ (u1_struct_0\ X0)) \\
& \quad (k1_pcomps_1\ X0\ X1)\ (k1_pcomps_1\ X0\ X2))\ (k1_pcomps_1\ X0\ (k7_subset_1 \\
& \quad (k1_zfmisc_1\ (u1_struct_0\ X0))\ X1\ X2))))
\end{aligned}$$