

t11_tops_1 (TMJTqGDVftp- MmWnb7GCmWQmTi6nsyHaEmyC)

October 27, 2020

Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v3_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \wedge (((v3_pre_topc X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0)))) \wedge ((v3_pre_topc X2 X0) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))))) \Rightarrow (v3_pre_topc (k3_xboole_0 X1 X2) X0) \quad (2)$$

Theorem 1

$$\forall X0. ((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (((v3_pre_topc X1 X0) \wedge (v3_pre_topc X2 X0)) \Rightarrow (v3_pre_topc (k9_subset_1 (u1_struct_0 X0) X1 X2) X0))))$$