

t11_uniform1 (TM-
PdQ2yhXLwuXtV8YE4WeZKnEM4h6ZSVjXA)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k18_complex1 : \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k17_complex1 : \iota \Rightarrow \iota$ be given. Let $k16_complex1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (\forall X1.(v1_xcmplx_0 X1) \Rightarrow (k17_complex1 (k6_xcmplx_0 X0 X1) = k17_complex1 (k6_xcmplx_0 X1 X0))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k18_complex1 X0 = k16_complex1 X0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k17_complex1 X0 = k16_complex1 X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k6_xcmplx_0 X0 X1)) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \quad (5)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (k18_complex1 (k6_xcmplx_0 X0 X1) = k18_complex1 (k6_xcmplx_0 X1 X0)))$$