

t122_pboole
(TMNnki7BBphXwZGr5LxAzaHwtNh33AJVYtS)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_reset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow (((k9_xtuple_0 X0 = k1_xboole_0) \vee (k10_xtuple_0 X0 = k1_xboole_0)) \Rightarrow (X0 = k1_xboole_0)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_reset_1 X0 X1 = k9_xtuple_0 X1) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (v1_partfun1 X1 X0) \Leftrightarrow (k1_reset_1 X0 X1 = X0) \quad (3)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k1_xboole_0) \wedge ((v1_funct_1 X0) \wedge (v1_partfun1 X0 k1_xboole_0)))) \Rightarrow (X0 = k1_xboole_0)$$