

## t122\_rvsum\_1

(TMSF1hjrwsuYZ4o4WmWAA3FUtLuWS73BV5Y)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v3\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k23\_rvsum\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_rvsum\_1 : \iota \Rightarrow \iota$  be given. Let  $k7\_binop\_2 : \iota \Rightarrow \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $k10\_rvsum\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k11\_binop\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xcmplx\_0 : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k30\_valued\_1 : \iota \Rightarrow \iota$  be given. Let  $k24\_valued\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_valued\_0 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge ((v3\_valued\_0 \\ & X0) \wedge (v1\_finseq\_1 X0)))) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 \\ & X1) \wedge ((v3\_valued\_0 X1) \wedge (v1\_finseq\_1 X1)))) \Rightarrow (\forall X2.(v1\_xreal\_0 \\ & X2) \Rightarrow ((k3\_finseq\_1 X0 = k3\_finseq\_1 X1) \Rightarrow (k23\_rvsum\_1 (k10\_rvsum\_1 \\ & X0 X2) X1 = k11\_binop\_2 X2 (k23\_rvsum\_1 X0 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow (k3\_xcmplx\_0 X0 (k4\_xcmplx\_0 np\_1) = k4\_xcmplx\_0 X0) \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge \\ & ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0) \Rightarrow (k7\_binop\_2 X0 = k4\_xcmplx\_0 X0) \quad (4)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge ((v3\_valued\_0 X0) \wedge (v1\_finseq\_1 X0)))) \Rightarrow (k6\_rvsum\_1 X0 = k30\_valued\_1 X0) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(k11\_binop\_2 X0 X1 = k3\_xcmplx\_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge((v3\_valued\_0 X0)\wedge(v1\_finseq\_1 X0))))\wedge(v1\_xreal\_0 X1))\Rightarrow(k10\_rvsum\_1 X0 X1 = k24\_valued\_1 X0 X1) \quad (7)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow((v1\_xcmplx\_0 (k4\_xcmplx\_0 X0))\wedge(v1\_xreal\_0 (k4\_xcmplx\_0 X0))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge((v3\_valued\_0 X0)\wedge(v1\_finseq\_1 X0))))\wedge((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge((v3\_valued\_0 X1)\wedge(v1\_finseq\_1 X1)))))\Rightarrow(m1\_subset\_1 (k23\_rvsum\_1 X0 X1) k1\_numbers) \quad (9)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_valued\_0 X0)))\Rightarrow(k30\_valued\_1 X0 = k24\_valued\_1 X0 (k4\_xcmplx\_0 np\_1)) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xreal\_0 X0)\wedge(v1\_xreal\_0 X1))\Rightarrow(k11\_binop\_2 X0 X1 = k11\_binop\_2 X1 X0) \quad (11)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0)\wedge(v3\_valued\_0 X0))\Rightarrow((v1\_relat\_1 X0)\wedge(v1\_valued\_0 X0)) \quad (12)$$

Assume the following.

$$\forall X0.(v1\_xreal\_0 X0)\Rightarrow(v1\_xcmplx\_0 X0) \quad (13)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k1\_numbers)\Rightarrow(v1\_xreal\_0 X0) \quad (14)$$

**Theorem 1**

$$\forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge((v3\_valued\_0 X0)\wedge(v1\_finseq\_1 X0))))\Rightarrow(\forall X1.((v1\_relat\_1 X1)\wedge((v1\_funct\_1 X1)\wedge((v3\_valued\_0 X1)\wedge(v1\_finseq\_1 X1))))\Rightarrow((k3\_finseq\_1 X0 = k3\_finseq\_1 X1)\Rightarrow(k23\_rvsum\_1 (k6\_rvsum\_1 X0) X1 = k7\_binop\_2 (k23\_rvsum\_1 X0 X1))))$$