

t123\_funct\_4 (TM-  
FQDzhT3YdvsvAscpUYFW3AiYATzpY3R2j)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_funct\_4 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (r1\_tarski X0 (k1\_funct\_4 X1 X0))) \quad (1)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (\forall X2.((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow (k1\_funct\_4 (k1\_funct\_4 X0 X1) X2 = k1\_funct\_4 X0 (k1\_funct\_4 X1 X2)))) \quad (2)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((r1\_tarski X0 X1) \Rightarrow (k1\_funct\_4 X1 X0 = X1))) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1.(((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \wedge ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1))) \Rightarrow ((v1\_relat\_1 (k1\_funct\_4 X0 X1)) \wedge (v1\_funct\_1 (k1\_funct\_4 X0 X1))) \quad (4)$$

**Theorem 1**

$$\forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1.((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (\forall X2.((v1\_relat\_1 X2) \wedge (v1\_funct\_1 X2)) \Rightarrow ((r1\_tarski X0 X1) \Rightarrow (r1\_tarski (k1\_funct\_4 X0 X2) (k1\_funct\_4 X1 X2))))))$$