

t124_relat_1 (TM-
PZdMY1aeiY1onm5f2vjWnFSfmBYHiQCb3)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (1)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (\forall X1. \forall X2. (X2 = k7_relat_1 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow (\exists X4. (k4_tarski X4 X3 \in X0) \wedge (X4 \in X1)))) \quad (2)$$

Theorem 1

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (\forall X2. (v1_relat_1 X2) \Rightarrow ((r1_tarski X1 X2) \Rightarrow (r1_tarski (k7_relat_1 X1 X0) (k7_relat_1 X2 X0))))$$