

t125_finseq_6

(TMYsaVQrPmg1scYrNyCWECzLH7hDCYhjPT8)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k17_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k8_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k12_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k9_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow ((X1 = k9_finseq_1 X0) \Leftrightarrow ((k3_finseq_1 X1 = np_1) \wedge (k10_xtuple_0 X1 = k1_tarski X0))) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\ (\forall X1. ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \Rightarrow (\forall X2. (v7_ordinal1 X2) \Rightarrow (k5_relat_1 (k7_finseq_1 \\ X0 X1) (k2_finseq_1 (k2_nat_1 (k3_finseq_1 X0) X2)) = k7_finseq_1 \\ X0 (k5_relat_1 X1 (k2_finseq_1 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0.k9_finseq_1 X0 = k5_finseq_1 X0 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_finseq_1 X1 X0)\wedge(m1_finseq_1 X2 X0))\Rightarrow(k8_finseq_1 X0 X1 X2 = k7_finseq_1 X1 X2) \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1 X1)\wedge(m1_finseq_1 X2 X0))\Rightarrow(k17_finseq_1 X0 X1 X2 = k16_finseq_1 X1 X2) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow(k12_finseq_1 X0 X1 = k5_finseq_1 X1) \quad (10)$$

Assume the following.

$$\forall X0.v1_finseq_1 (k5_finseq_1 X0) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(v7_ordinal1 (k2_xcmplx_0 X0 X1)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (14)$$

Assume the following.

$$\forall X0.(v1_relat_1 (k9_finseq_1 X0)\wedge(v1_funct_1 (k9_finseq_1 X0))) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_finseq_1 X1 X0)\wedge(m1_finseq_1 X2 X0))\Rightarrow(m2_finseq_1 (k8_finseq_1 X0 X1 X2) X0) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1 X1)\wedge(m1_finseq_1 X2 X0))\Rightarrow(m2_finseq_1 (k17_finseq_1 X0 X1 X2) X0) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(m1_subset_1 X1 X0))\Rightarrow(m2_finseq_1 (k12_finseq_1 X0 X1) X0) \quad (18)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(\forall X1.((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1)))\Rightarrow(k16_finseq_1 X0 X1 = k5_relat_1 X1 (k2_finseq_1 X0))) \quad (19)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_nat_1 X1 X0) \quad (20)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (21)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(m2_finseq_1 X1 X0)\Rightarrow(\forall X2.(m1_subset_1 X2 k5_numbers)\Rightarrow(\forall X3.(m1_subset_1 X3 X0)\Rightarrow(k17_finseq_1 X0 (k2_nat_1 X2 np_1) (k8_finseq_1 X0 (k12_finseq_1 X0 X3) X1) = k8_finseq_1 X0 (k12_finseq_1 X0 X3) (k17_finseq_1 X0 X2 X1))))))$$