

# t128\_funct\_7 (TMUAajjTSZAih- WhTrXDQ8jn9KkTP9DXbzRd)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_funct\_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_relat\_1 X0) \wedge (v1\_funct\_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ & (X2 \in k9\_xtuple\_0 X0) \Rightarrow (k1\_funct\_1 (k2\_funct\_7 X0 X2 X1) X2 = X1)) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 X1) \Rightarrow ((v1\_xboole\_0 X1) \vee (X0 \in X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (v1\_partfun1 X1 X0 \Leftrightarrow (k1\_relset\_1 X0 X1 = X0)) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (\neg v1\_xboole\_0 X1) \Rightarrow (\forall X2. (m1\_subset\_1 \\ & X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v1\_funct\_2 X2 X0 X1) \Rightarrow ( \\ & v1\_partfun1 X2 X0))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v4\_relat\_1 X2 X0) \wedge (v5\_relat\_1 X2 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (7)$$

**Theorem 1**

$$\begin{aligned} &\forall X0.(\neg v1\_xboole\_0 X0)\Rightarrow(\forall X1.(\neg v1\_xboole\_0 X1)\Rightarrow \\ &(\forall X2.((v1\_funct\_1 X2)\wedge((v1\_funct\_2 X2 X0 X1)\wedge(m1\_subset\_1 \\ &X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))))\Rightarrow(\forall X3.(m1\_subset\_1 \\ &X3 X0)\Rightarrow(\forall X4.k1\_funct\_1 (k2\_funct\_7 X2 X3 X4) X3 = X4)))) \end{aligned}$$