

t12_anproj_2

(TMa5WjAhriFor5zjNYBahWPFq5FQ8vRSMdb)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k9_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funcsdom : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_funcsdom : \iota \Rightarrow \iota$ be given. Let $k2_funcsdom : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_funcsdom : \iota \Rightarrow \iota$ be given. Let $k1_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_funcsdom : \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0. (\neg v1_xboole_0 X0) \Rightarrow (\forall X1. (m2_funct_2 X1 X0 k1_numbers \\
 & (k9_funct_2 X0 k1_numbers)) \Rightarrow (\forall X2. (m2_funct_2 X2 X0 k1_numbers \\
 & (k9_funct_2 X0 k1_numbers)) \Rightarrow (\forall X3. (m2_funct_2 X3 X0 k1_numbers \\
 & (k9_funct_2 X0 k1_numbers)) \Rightarrow (\forall X4. (m1_subset_1 X4 X0) \Rightarrow \\
 & (\forall X5. (m1_subset_1 X5 X0) \Rightarrow (\forall X6. (m1_subset_1 X6 X0) \Rightarrow \\
 & (((k3_funct_2 X0 k1_numbers X1 X4 = np_1) \wedge ((\forall X7. (X7 \in X0) \Rightarrow \\
 & ((X7 = X4) \vee (k1_funct_1 X1 X7 = k6_numbers)))) \wedge ((k3_funct_2 X0 k1_numbers \\
 & X2 X5 = np_1) \wedge ((\forall X7. (X7 \in X0) \Rightarrow ((X7 = X5) \vee (k1_funct_1 X2 X7 = \\
 & k6_numbers)))) \wedge ((k3_funct_2 X0 k1_numbers X3 X6 = np_1) \wedge (\forall X7. \\
 & (X7 \in X0) \Rightarrow ((X7 = X6) \vee (k1_funct_1 X3 X7 = k6_numbers)))))) \Rightarrow ((X4 = \\
 & X5) \vee ((X4 = X6) \vee ((X5 = X6) \vee (\forall X7. (m1_subset_1 X7 k1_numbers) \Rightarrow \\
 & (\forall X8. (m1_subset_1 X8 k1_numbers) \Rightarrow (\forall X9. (m1_subset_1 \\
 & X9 k1_numbers) \Rightarrow ((r2_funct_2 X0 k1_numbers (k1_funcsdom X0 k1_numbers \\
 & (k5_funcsdom X0) (k1_funcsdom X0 k1_numbers (k5_funcsdom X0) (\\
 & k2_funcsdom X0 k1_numbers k1_numbers (k9_funct_2 X0 k1_numbers) \\
 & (k7_funcsdom X0) (k1_domain_1 k1_numbers (k9_funct_2 X0 k1_numbers) \\
 & X7 X1)) (k2_funcsdom X0 k1_numbers k1_numbers (k9_funct_2 X0 k1_numbers) \\
 & (k7_funcsdom X0) (k1_domain_1 k1_numbers (k9_funct_2 X0 k1_numbers) \\
 & X8 X2))) (k2_funcsdom X0 k1_numbers k1_numbers (k9_funct_2 X0 k1_numbers) \\
 & (k7_funcsdom X0) (k1_domain_1 k1_numbers (k9_funct_2 X0 k1_numbers) \\
 & X9 X3))) (k8_funcsdom X0) \Rightarrow ((X7 = k6_numbers) \wedge ((X8 = k6_numbers) \wedge \\
 & (X9 = k6_numbers))))))))))))))
 \end{aligned}$$

(1)

Assume the following.

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow \\
& (\exists X2.(m2_funct_2 X2 X0 k1_numbers (k9_funct_2 X0 k1_numbers)) \wedge \\
& ((k3_funct_2 X0 k1_numbers X2 X1 = np_1) \wedge (\forall X3.(X3 \in X0) \Rightarrow \\
& ((X3 = X1) \vee (k1_funct_1 X2 X3 = k6_numbers))))))
\end{aligned} \tag{2}$$

Theorem 1

$$\begin{aligned}
& \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow \\
& (\forall X2.(m1_subset_1 X2 X0) \Rightarrow (\forall X3.(m1_subset_1 X3 X0) \Rightarrow \\
& (\neg (X1 \neq X2) \wedge (X1 \neq X3) \wedge (X2 \neq X3) \wedge (\forall X4.(m2_funct_2 X4 X0 k1_numbers \\
& (k9_funct_2 X0 k1_numbers)) \Rightarrow (\forall X5.(m2_funct_2 X5 X0 k1_numbers \\
& (k9_funct_2 X0 k1_numbers)) \Rightarrow (\forall X6.(m2_funct_2 X6 X0 k1_numbers \\
& (k9_funct_2 X0 k1_numbers)) \Rightarrow (\exists X7.(m1_subset_1 X7 k1_numbers) \wedge \\
& (\exists X8.(m1_subset_1 X8 k1_numbers) \wedge (\exists X9.(m1_subset_1 \\
& X9 k1_numbers) \wedge ((r2_funct_2 X0 k1_numbers (k1_funcsdom X0 k1_numbers \\
& (k5_funcsdom X0) (k1_funcsdom X0 k1_numbers (k5_funcsdom X0) (\\
& k2_funcsdom X0 k1_numbers k1_numbers (k9_funct_2 X0 k1_numbers) \\
& (k7_funcsdom X0) (k1_domain_1 k1_numbers (k9_funct_2 X0 k1_numbers) \\
& X7 X4) (k2_funcsdom X0 k1_numbers k1_numbers (k9_funct_2 X0 k1_numbers) \\
& (k7_funcsdom X0) (k1_domain_1 k1_numbers (k9_funct_2 X0 k1_numbers) \\
& X8 X5)) (k2_funcsdom X0 k1_numbers k1_numbers (k9_funct_2 X0 k1_numbers) \\
& (k7_funcsdom X0) (k1_domain_1 k1_numbers (k9_funct_2 X0 k1_numbers) \\
& X9 X6))) (k8_funcsdom X0) \wedge (\neg (X7 = k6_numbers) \wedge ((X8 = k6_numbers) \wedge \\
& (X9 = k6_numbers))))))))))))))
\end{aligned}$$