

## t12\_fib\_num2

(TMcF1QHgzL9ChBcqgfs7Ss1tt8YHvSPXCKD)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $r1\_nat\_d : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7\_ordinal1 X2) \Rightarrow ((r1\_nat\_d X0 X1) \Rightarrow (r1\_nat\_d X0 (k3\_xcmplx\_0 \\ & X1 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7\_ordinal1 X2) \Rightarrow (((r1\_nat\_d X0 X1) \wedge (r1\_nat\_d X0 X2)) \Rightarrow (r1\_nat\_d \\ & X0 (k2\_xcmplx\_0 X1 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\ & X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers) \wedge (v7\_ordinal1 X1)) \Rightarrow (k4\_nat\_1 X0 X1 = k3\_xcmplx\_0 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0) \wedge (m1\_subset\_1 X1 k5\_numbers)) \Rightarrow (k3\_nat\_1 X0 X1 = k3\_xcmplx\_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(k2\_nat\_1 X0 X1 = k2\_xcmplx\_0 X0 X1) \quad (7)$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1)\wedge(v3\_ordinal1 k4\_ordinal1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v7\_ordinal1 X0)\wedge(v7\_ordinal1 X1))\Rightarrow(v7\_ordinal1 (k3\_xcmplx\_0 X0 X1)) \quad (9)$$

Assume the following.

$$\neg v1\_xboole\_0 k1\_numbers \quad (10)$$

Assume the following.

$$m1\_subset\_1 k5\_numbers (k1\_zfmisc\_1 k1\_numbers) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k5\_numbers)\wedge(v7\_ordinal1 X1))\Rightarrow(m2\_subset\_1 (k4\_nat\_1 X0 X1) k1\_numbers k5\_numbers) \quad (12)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (13)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(v7\_ordinal1 X0)\Rightarrow(\forall X1.(m2\_subset\_1 X1 k1\_numbers \\ & k5\_numbers)\Rightarrow(\forall X2.(m2\_subset\_1 X2 k1\_numbers k5\_numbers)\Rightarrow \\ & (\forall X3.(m2\_subset\_1 X3 k1\_numbers k5\_numbers)\Rightarrow(\forall X4. \\ & (m2\_subset\_1 X4 k1\_numbers k5\_numbers)\Rightarrow(((r1\_nat\_d X1 X2)\wedge(r1\_nat\_d \\ & X1 X0))\Rightarrow(r1\_nat\_d X1 (k2\_nat\_1 (k4\_nat\_1 X2 X3) (k3\_nat\_1 X0 X4))))))))) \end{aligned}$$