

t12_funct_1
(TMPNj6rEia1FpETgJxHdmw3ABnkznA5geN2)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \wedge ((v1_relat_1 X1) \wedge (v1_funct_1 X1))) \Rightarrow ((v1_relat_1 (k3_relat_1 X0 X1)) \wedge (v1_funct_1 (k3_relat_1 X0 X1))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. v1_relat_1 (k3_relat_1 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (v1_relat_1 X2) \Rightarrow ((X2 = k3_relat_1 X0 X1) \Leftrightarrow (\forall X3. \forall X4. (k4_tarski X3 X4 \in X2) \Leftrightarrow (\exists X5. (k4_tarski X3 X5 \in X0) \wedge (k4_tarski X5 X4 \in X1)))) \quad (3)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. ((X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (k4_tarski X1 X2 \in X0))) \wedge ((\neg X1 \in k9_xtuple_0 X0) \Rightarrow ((X2 = k1_funct_1 X0 X1) \Leftrightarrow (X2 = k1_xboole_0)))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k9_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. k4_tarski X2 X3 \in X0)) \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 (k3_relat_1 X2 X1)) \Rightarrow (k1_funct_1 (k3_relat_1 X2 X1) X0 = k1_funct_1 X1 (k1_funct_1 X2 X0))))$$