

# t12\_funct\_2 (TMM- nGQDypa9Au17mC6MYR836vL8Q8GeLP38)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v1\_funct\_1 X2) \wedge (m1\_subset\_1 \\ & \quad X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))) \Rightarrow (\forall X3. ((v1\_funct\_1 \\ & \quad X3) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))) \Rightarrow (((k1\_relset\_1 \\ & \quad X0 X2 = k1\_relset\_1 X0 X3) \wedge (\forall X4. (m1\_subset\_1 X4 X0) \Rightarrow ((X4 \in \\ & \quad k1\_relset\_1 X0 X2) \Rightarrow (k1\_funct\_1 X2 X4 = k1\_funct\_1 X3 X4)))) \Rightarrow (X2 = \\ & \quad X3))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((m1\_subset\_1 X2 \\ & \quad (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ & \quad (k2\_zfmisc\_1 X0 X1)))) \Rightarrow ((r2\_relset\_1 X0 X1 X2 X3) \Rightarrow (r2\_relset\_1 \\ & \quad X0 X1 X3 X2)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. ((m1\_subset\_1 X2 \\ & \quad (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\ & \quad (k2\_zfmisc\_1 X0 X1)))) \Rightarrow ((r2\_relset\_1 X0 X1 X2 X3) \Leftrightarrow (X2 = X3)) \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & \quad (k2\_zfmisc\_1 X0 X1))) \Rightarrow (((X1 \neq k1\_xboole\_0) \Rightarrow ((v1\_funct\_2 X2 X0 \\ & \quad X1) \Leftrightarrow (X0 = k1\_relset\_1 X0 X2))) \wedge ((X1 = k1\_xboole\_0) \Rightarrow ((v1\_funct\_2 \\ & \quad X2 X0 X1) \Leftrightarrow (X2 = k1\_xboole\_0)))) \end{aligned} \tag{4}$$

**Theorem 1**

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v1\_funct\_1 X2) \wedge ((v1\_funct\_2 \\ & X2 X0 X1) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))))) \Rightarrow \\ & (\forall X3. ((v1\_funct\_1 X3) \wedge ((v1\_funct\_2 X3 X0 X1) \wedge (m1\_subset\_1 \\ & X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))))) \Rightarrow ((\forall X4. (X4 \in X0) \Rightarrow \\ & (k1\_funct\_1 X2 X4 = k1\_funct\_1 X3 X4)) \Rightarrow (r2\_reset\_1 X0 X1 X2 X3))) \end{aligned}$$