

t12_latsum_1

(TMRx7ACaCc6mmWewNZD6i2NRdAYGhCrdXGX)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. (X2 = k3_xboole_0 X0 X1) &\Leftrightarrow (\forall X3. \\ (X3 \in X2) &\Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} \forall X0. ((\neg v2_struct_0 X0) \wedge (l1_orders_2 X0)) &\Rightarrow (\forall X1. \\ ((\neg v2_struct_0 X1) \wedge (l1_orders_2 X1)) &\Rightarrow (\forall X2. (X2 \in k3_xboole_0 \\ (u1_struct_0 X0) (u1_struct_0 X1)) &\Rightarrow (m1_subset_1 X2 (u1_struct_0 \\ X0)))) \end{aligned}$$