

t12_pcomps_1 (TMQHtBXe- HVx5MVUXrwgEtRvmjN4nS81S1MM)

October 27, 2020

Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k1_pcomps_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k2_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \neg (X0 \in X1) \wedge (v1_xboole_0 X1) \quad (1)$$

Assume the following.

$$v1_xboole_0 k1_xboole_0 \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. (l1_pre_topc X0) \Rightarrow (\forall X1. (m1_subset_1 X1 (k1_zfmisc_1 \\ (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow (\forall X2. (m1_subset_1 X2 \\ (k1_zfmisc_1 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow ((X2 = k1_pcomps_1 \\ X0 X1) \Leftrightarrow (\forall X3. (m1_subset_1 X3 (k1_zfmisc_1 (u1_struct_0 \\ X0)))) \Rightarrow ((X3 \in X2) \Leftrightarrow (\exists X4. (m1_subset_1 X4 (k1_zfmisc_1 (u1_struct_0 \\ X0)))) \wedge ((X3 = k2_pre_topc X0 X4) \wedge (X4 \in X1))))))))) \end{aligned} \quad (3)$$

Theorem 1

$$\begin{aligned} \forall X0. ((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. \\ (m1_subset_1 X1 (k1_zfmisc_1 (k1_zfmisc_1 (u1_struct_0 X0)))) \Rightarrow \\ ((X1 = k1_xboole_0) \Rightarrow (k1_pcomps_1 X0 X1 = k1_xboole_0))) \end{aligned}$$