

t12_seq_1

(TMVsm5DQJTHdY7kN3xFsns4z8vDJU4Rku79)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k56_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k54_valued_1 : \iota \Rightarrow \iota$ be given. Let $k17_complex1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k16_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \neg(v1_xboole_0 X0) \wedge ((X0 \neq X1) \wedge (v1_xboole_0 X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (X0 \in X1) \Rightarrow (m1_subset_1 X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow (\forall X1. k1_funct_1 (k54_valued_1 X0) X1 = k17_complex1 (k1_funct_1 X0 X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (((v1_funct_1 X2) \wedge ((v1_funct_2 X2 X0 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \wedge ((v1_funct_1 X3) \wedge ((v1_funct_2 X3 X0 X1) \wedge (m1_subset_1 X3 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow ((r2_funct_2 X0 X1 X2 X3) \Leftrightarrow (X2 = X3)) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Leftrightarrow(m1_subset_1 X2 X1)) \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_membered X1)\wedge((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))\Rightarrow(k56_valued_1 X0 X1 X2 = k54_valued_1 X2) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v3_valued_0 X0)))\Rightarrow(k1_seq_1 X0 X1 = k1_funct_1 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0)\Rightarrow(k18_complex1 X0 = k16_complex1 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0)\Rightarrow(k17_complex1 X0 = k16_complex1 X0) \quad (11)$$

Assume the following.

$$\exists X0.v1_xboole_0 X0 \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(((\neg v1_xboole_0 X1)\wedge(v1_membered X1))\wedge((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))))\Rightarrow((v1_funct_1 (k54_valued_1 X2))\wedge(v1_partfun1 (k54_valued_1 X2) X0)) \quad (13)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 X0)))\Rightarrow(v1_xcmplx_0 (k1_funct_1 X0 X1)) \quad (15)$$

Assume the following.

$$v3_membered\ k1_numbers \quad (16)$$

Assume the following.

$$v1_xboole_0\ k1_xboole_0 \quad (17)$$

Assume the following.

$$\neg v1_xboole_0\ k1_numbers \quad (18)$$

Assume the following.

$$m1_subset_1\ k5_numbers\ (k1_zfmisc_1\ k1_numbers) \quad (19)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((v1_membered\ X1)\wedge((v1_funct_1 \\ X2)\wedge(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1))))\Rightarrow((v1_funct_1 \\ (k56_valued_1\ X0\ X1\ X2))\wedge(m1_subset_1\ (k56_valued_1\ X0\ X1\ X2)\ (\\ k1_zfmisc_1\ (k2_zfmisc_1\ X0\ k1_numbers)))) \end{aligned} \quad (20)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_valued_0\ X0)))\Rightarrow \\ ((v1_relat_1\ (k54_valued_1\ X0))\wedge((v1_funct_1\ (k54_valued_1 \\ X0))\wedge(v3_valued_0\ (k54_valued_1\ X0)))) \end{aligned} \quad (21)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1 \\ (k2_zfmisc_1\ X0\ X1)))\Rightarrow(((X1\neq k1_xboole_0)\Rightarrow((v1_funct_2\ X2\ X0 \\ X1)\Leftrightarrow(X0 = k1_relset_1\ X0\ X2)))\wedge((X1 = k1_xboole_0)\Rightarrow((v1_funct_2 \\ X2\ X0\ X1)\Leftrightarrow(X2 = k1_xboole_0)))) \end{aligned} \quad (22)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_valued_0\ X0)))\Rightarrow \\ (\forall X1.((v1_relat_1\ X1)\wedge((v1_funct_1\ X1)\wedge(v3_valued_0 \\ X1)))\Rightarrow((X1 = k54_valued_1\ X0)\Leftrightarrow((k9_xtuple_0\ X1 = k9_xtuple_0\ X0)\wedge \\ (\forall X2.(X2 \in k9_xtuple_0\ X1)\Rightarrow(k1_funct_1\ X1\ X2 = k17_complex1 \\ (k1_funct_1\ X0\ X2)))))) \end{aligned} \quad (23)$$

Assume the following.

$$\forall X0.(v3_membered\ X0)\Rightarrow(v1_membered\ X0) \quad (24)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1 \\ (k2_zfmisc_1\ X0\ X1)))\Rightarrow((v4_relat_1\ X2\ X0)\wedge(v5_relat_1\ X2\ X1)) \end{aligned} \quad (25)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (26)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v1_partfun1 X2 X0)\Rightarrow(v1_funct_2 X2 X0 X1)) \quad (27)$$

Assume the following.

$$\forall X0.\forall X1.(v3_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v3_valued_0 X2)) \quad (28)$$

Assume the following.

$$\forall X0.\forall X1.(v1_membered X1)\Rightarrow(\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_valued_0 X2)) \quad (29)$$

Theorem 1

$$\begin{aligned} & \forall X0.((v1_funct_1 X0)\wedge((v1_funct_2 X0 k5_numbers k1_numbers)\wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers))))))\Rightarrow \\ & (\forall X1.((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers k1_numbers)\wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k1_numbers))))))\Rightarrow \\ & ((r2_funct_2 k5_numbers k1_numbers X0 (k56_valued_1 k5_numbers \\ & k1_numbers X1))\Leftrightarrow(\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers)\Rightarrow \\ & (k1_seq_1 X0 X2 = k18_complex1 (k1_seq_1 X1 X2)))))) \end{aligned}$$