

t12_sheffer2

(TMTr3QLWw9ABF9KPqNUK7oxSEDb9VYRm1dK)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v1_sheffer2 : \iota \Rightarrow o$ be given. Let $l1_sheffer1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k5_sheffer1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & \quad X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 (k5_sheffer1 \\ & \quad X0 X1 X1) (k5_sheffer1 X0 X2 X1) = X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(((\neg v2_struct_0 X0) \wedge (l1_sheffer1 \\ & X0)) \wedge ((m1_subset_1 X1 (u1_struct_0 X0)) \wedge (m1_subset_1 X2 (u1_struct_0 \\ & \quad X0)))) \Rightarrow (m1_subset_1 (k5_sheffer1 X0 X1 X2) (u1_struct_0 X0)) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_sheffer2 X0) \wedge (l1_sheffer1 \\ & \quad X0))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2. \\ & (m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (k5_sheffer1 X0 X1 (k5_sheffer1 \\ & \quad X0 X2 (k5_sheffer1 X0 X1 X1)) = k5_sheffer1 X0 X1 X1))) \end{aligned}$$