

t12\_sppol\_2 (TMSS-  
ByRy5LZD5aTX5uQUVP1oASZPubKXQdd)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k15\_euclid : \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k3\_topreal1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k12\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v5\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $l1\_struct\_0 : \iota \Rightarrow o$  be given. Let  $l1\_rltopsp1 : \iota \Rightarrow o$  be given. Let  $l1\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v1\_funct\_1 X1) \wedge (v1\_finseq\_1 \\ X1))) \Rightarrow ((X1 = k9\_finseq\_1 X0) \Leftrightarrow ((k3\_finseq\_1 X1 = np\_1) \wedge (k10\_xtuple\_0 \\ X1 = k1\_tarski X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. (v7\_ordinal1 X0) \Rightarrow (\forall X1. (m2\_finseq\_1 X1 (u1\_struct\_0 \\ (k15\_euclid X0))) \Rightarrow (((k3\_finseq\_1 X1 = k6\_numbers) \vee (k3\_finseq\_1 \\ X1 = np\_1)) \Leftrightarrow (k3\_topreal1 X0 X1 = k1\_xboole\_0))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} ((v2\_xxreal\_0 np\_2) \wedge (m2\_subset\_1 np\_2 k1\_numbers k5\_numbers)) \wedge \\ ((m1\_subset\_1 np\_2 k5\_numbers) \wedge (m1\_subset\_1 np\_2 k1\_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(m1\_subset\_1 X1 X0))\Rightarrow (k12\_finseq\_1 X0 X1 = k5\_finseq\_1 X1) \quad (6)$$

Assume the following.

$$\forall X0.v1\_finseq\_1 (k5\_finseq\_1 X0) \quad (7)$$

Assume the following.

$$\forall X0.(v1\_relat\_1 (k5\_finseq\_1 X0))\wedge(v1\_funct\_1 (k5\_finseq\_1 X0)) \quad (8)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow((\neg v2\_struct\_0 (k15\_euclid X0))\wedge (v5\_rltopsp1 (k15\_euclid X0))) \quad (9)$$

Assume the following.

$$\forall X0.((\neg v2\_struct\_0 X0)\wedge(l1\_struct\_0 X0))\Rightarrow(\neg v1\_xboole\_0 (u1\_struct\_0 X0)) \quad (10)$$

Assume the following.

$$\forall X0.(l1\_rltopsp1 X0)\Rightarrow((l1\_rlvect\_1 X0)\wedge(l1\_pre\_topc X0)) \quad (11)$$

Assume the following.

$$\forall X0.(l1\_pre\_topc X0)\Rightarrow(l1\_struct\_0 X0) \quad (12)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0)\Rightarrow((v5\_rltopsp1 (k15\_euclid X0))\wedge (l1\_rltopsp1 (k15\_euclid X0))) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1\_xboole\_0 X0)\wedge(m1\_subset\_1 X1 X0))\Rightarrow (m2\_finseq\_1 (k12\_finseq\_1 X0 X1) X0) \quad (14)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1)\Rightarrow(v7\_ordinal1 X0) \quad (15)$$

**Theorem 1**

$$\forall X0.(m1\_subset\_1 X0 (u1\_struct\_0 (k15\_euclid np\_2)))\Rightarrow (k3\_topreal1 np\_2 (k12\_finseq\_1 (u1\_struct\_0 (k15\_euclid np\_2)) X0) = k1\_xboole\_0)$$