

t12_waybel_8

(TMMkk34VTKKraccGFUYAER6q2tZrBuM815J)

October 27, 2020

Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $g1_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $u1_orders_2 : \iota \Rightarrow \iota$ be given. Let $v3_orders_2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_relat_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ X0 X0))) \Rightarrow (\forall X2. \forall X3. (g1_orders_2 X0 X1 = g1_orders_2 \\ X2 X3) \Rightarrow ((X0 = X2) \wedge (X1 = X3))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. (l1_orders_2 X0) \Rightarrow (m1_subset_1 (u1_orders_2 X0) (k1_zfmisc_1 \\ (k2_zfmisc_1 (u1_struct_0 X0) (u1_struct_0 X0)))) \quad (2)$$

Assume the following.

$$\forall X0. (l1_orders_2 X0) \Rightarrow ((v3_orders_2 X0) \Leftrightarrow (r1_relat_2 (\\ u1_orders_2 X0) (u1_struct_0 X0))) \quad (3)$$

Theorem 1

$$\begin{aligned} \forall X0. (l1_orders_2 X0) \Rightarrow (\forall X1. (l1_orders_2 X1) \Rightarrow ((\\ (g1_orders_2 (u1_struct_0 X0) (u1_orders_2 X0) = g1_orders_2 (\\ u1_struct_0 X1) (u1_orders_2 X1)) \wedge (v3_orders_2 X0)) \Rightarrow (v3_orders_2 \\ X1))) \end{aligned}$$