

t12_xtuple_0
(TMX5U81kaM3VtVPRpBFU1QMa6s7jxgWxD13)

October 27, 2020

Let $k11_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k3_xtuple_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. k3_xtuple_0 X0 X1 X2 = k4_tarski (k4_tarski X0 X1) X2 \quad (1)$$

Assume the following.

$$\forall X0. k11_xtuple_0 X0 = k9_xtuple_0 (k9_xtuple_0 X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k9_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. k4_tarski X2 X3 \in X0)) \quad (3)$$

Theorem 1

$$\forall X0. \forall X1. \neg (X0 \in k11_xtuple_0 X1) \wedge (\forall X2. \forall X3. \neg k3_xtuple_0 X0 X2 X3 \in X1)$$