

t12_xxreal_3
(TMQzE7s9dAJyHXg5hEsz9xw7Scr3h8Zsx8i)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k3_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_3 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2.(v1_xxreal_0 X2) \Rightarrow ((k1_xxreal_3 X0 X1 = k1_xxreal_3 X0 X2) \Rightarrow (X1 = X2)))) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (k2_xxreal_3 (k2_xxreal_3 X0) = X0) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (v1_xxreal_0 (k2_xxreal_3 X0)) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (k3_xxreal_3 X0 X1 = k1_xxreal_3 X0 (k2_xxreal_3 X1))) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xxreal_0 X0) \quad (5)$$

Theorem 1

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2.(v1_xxreal_0 X2) \Rightarrow ((k3_xxreal_3 X0 X1 = k3_xxreal_3 X0 X2) \Rightarrow (X1 = X2))))$$