

t130\_relat\_1  
(TMWikKBv2WgAg6RaQPvmydaTi7SPGLsn2c1)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k7\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_relat\_1 : \iota \Rightarrow \iota$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_relat\_1 X0) \Rightarrow (v1\_relat\_1 (k2\_relat\_1 X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_relat\_1 X0) \Rightarrow (\forall X1.(v1\_relat\_1 X1) \Rightarrow ((X1 = k2\_relat\_1 X0) \Leftrightarrow (\forall X2.\forall X3.(k4\_tarski X2 X3 \in X1) \Leftrightarrow (k4\_tarski X3 X2 \in X0)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(X2 = k3\_xboole\_0 X0 X1) \Leftrightarrow (\forall X3.(X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(r1\_tarski X0 X1) \Leftrightarrow (\forall X2.(X2 \in X0) \Rightarrow (X2 \in X1)) \quad (4)$$

Assume the following.

$$\forall X0.(v1\_relat\_1 X0) \Rightarrow (\forall X1.\forall X2.(X2 = k7\_relat\_1 X0 X1) \Leftrightarrow (\forall X3.(X3 \in X2) \Leftrightarrow (\exists X4.(k4\_tarski X4 X3 \in X0) \wedge (X4 \in X1)))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k9\_xtuple\_0 X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.k4\_tarski X2 X3 \in X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.k3\_xboole\_0 X0 X1 = k3\_xboole\_0 X1 X0 \quad (7)$$

**Theorem 1**

$$\forall X0.\forall X1.(v1\_relat\_1 X1)\Rightarrow(r1\_tarSKI (k3\_xboole\_0 (k9\_xtuple\_0 X1) X0) (k7\_relat\_1 (k2\_relat\_1 X1) (k7\_relat\_1 X1 X0)))$$