

t132_zf_lang1

(TMMK8kR8SxNppeHLUnY4DBebtuSzSZbW8qf)

October 27, 2020

Let $v1_zf_lang : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zf_lang : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $r2_zf_model : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k11_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_zf_lang : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & (m2_subset_1 X1 k5_numbers k1_zf_lang) \Rightarrow (\forall X2. (\neg v1_xboole_0 \\ & X2) \Rightarrow (r2_zf_model X2 (k11_zf_lang (k8_zf_lang X1 X0) X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_zf_lang X0) \wedge (m2_finseq_1 X0 k5_numbers)) \Rightarrow (\forall X1. \\ & ((v1_zf_lang X1) \wedge (m2_finseq_1 X1 k5_numbers)) \Rightarrow (\forall X2. (\\ & (v1_zf_lang X2) \wedge (m2_finseq_1 X2 k5_numbers)) \Rightarrow (\forall X3. (\neg \\ & v1_xboole_0 X3) \Rightarrow (((r2_zf_model X3 (k11_zf_lang X0 X1)) \wedge (r2_zf_model \\ & X3 (k11_zf_lang X1 X2))) \Rightarrow (r2_zf_model X3 (k11_zf_lang X0 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((m1_subset_1 X0 k1_zf_lang) \wedge ((v1_zf_lang \\ & X1) \wedge (m1_finseq_1 X1 k5_numbers))) \Rightarrow (v1_zf_lang (k8_zf_lang X0 \\ & X1)) \end{aligned} \quad (5)$$

Assume the following.

$$\neg v1_xboole_0 \ k1_zf_lang \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 \ X0 \ k1_zf_lang)\wedge(m1_finseq_1 \ X1 \ k5_numbers))\Rightarrow(m2_finseq_1 \ (k8_zf_lang \ X0 \ X1) \ k5_numbers) \quad (7)$$

Assume the following.

$$m1_subset_1 \ k1_zf_lang \ (k1_zfmisc_1 \ k5_numbers) \quad (8)$$

Assume the following.

$$\forall X0.(v1_xboole_0 \ X0)\Rightarrow(\forall X1.(m1_subset_1 \ X1 \ (k1_zfmisc_1 \ X0))\Rightarrow(v1_xboole_0 \ X1)) \quad (9)$$

Theorem 1

$$\begin{aligned} &\forall X0.((v1_zf_lang \ X0)\wedge(m2_finseq_1 \ X0 \ k5_numbers))\Rightarrow(\forall X1. \\ &\quad ((v1_zf_lang \ X1)\wedge(m2_finseq_1 \ X1 \ k5_numbers))\Rightarrow(\forall X2.(\\ &\quad m2_subset_1 \ X2 \ k5_numbers \ k1_zf_lang)\Rightarrow(\forall X3.(\neg v1_xboole_0 \\ X3)\Rightarrow((r2_zf_model \ X3 \ (k11_zf_lang \ X0 \ (k8_zf_lang \ X2 \ X1)))\Rightarrow(r2_zf_model \\ &\quad X3 \ (k11_zf_lang \ X0 \ X1)))))) \end{aligned}$$