

# t133\_zf\_lang1 (TMK Ler- DeNm v1MsDn2GMp9WQnUZu8aEjTgsy)

October 27, 2020

Let  $v1\_zf\_lang : \iota \Rightarrow o$  be given. Let  $m2\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zf\_lang : \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $r2\_zf\_model : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k11\_zf\_lang : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k13\_zf\_lang : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zf\_misc\_1 : \iota \Rightarrow \iota$  be given. Let  $m1\_finseq\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_zf\_lang X0) \wedge (m2\_finseq\_1 X0 k5\_numbers)) \Rightarrow (\forall X1. \\ & (m2\_subset\_1 X1 k5\_numbers k1\_zf\_lang) \Rightarrow (\forall X2.(\neg v1\_xboole\_0 \\ & X2) \Rightarrow (r2\_zf\_model X2 (k11\_zf\_lang X0 (k13\_zf\_lang X1 X0)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1\_zf\_lang X0) \wedge (m2\_finseq\_1 X0 k5\_numbers)) \Rightarrow (\forall X1. \\ & ((v1\_zf\_lang X1) \wedge (m2\_finseq\_1 X1 k5\_numbers)) \Rightarrow (\forall X2.( \\ & (v1\_zf\_lang X2) \wedge (m2\_finseq\_1 X2 k5\_numbers)) \Rightarrow (\forall X3.(\neg \\ & v1\_xboole\_0 X3) \Rightarrow (((r2\_zf\_model X3 (k11\_zf\_lang X0 X1)) \wedge (r2\_zf\_model \\ & X3 (k11\_zf\_lang X1 X2))) \Rightarrow (r2\_zf\_model X3 (k11\_zf\_lang X0 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ & (m1\_subset\_1 X1 (k1\_zf\_misc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\ & X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (m2\_finseq\_1 X1 X0) \Leftrightarrow (m1\_finseq\_1 X1 X0) \quad (4)$$

Assume the following.

$$\neg v1\_xboole\_0 k1\_zf\_lang \quad (5)$$

Assume the following.

$$m1\_subset\_1 k1\_zf\_lang (k1\_zf\_misc\_1 k5\_numbers) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((m1\_subset\_1 X0 k1\_zf\_lang)\wedge((v1\_zf\_lang X1)\wedge(m1\_finseq\_1 X1 k5\_numbers)))\Rightarrow((v1\_zf\_lang (k13\_zf\_lang X0 X1))\wedge(m2\_finseq\_1 (k13\_zf\_lang X0 X1) k5\_numbers)) \quad (7)$$

Assume the following.

$$\forall X0.(v1\_xboole\_0 X0)\Rightarrow(\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 X0))\Rightarrow(v1\_xboole\_0 X1)) \quad (8)$$

**Theorem 1**

$$\forall X0.((v1\_zf\_lang X0)\wedge(m2\_finseq\_1 X0 k5\_numbers))\Rightarrow(\forall X1.((v1\_zf\_lang X1)\wedge(m2\_finseq\_1 X1 k5\_numbers))\Rightarrow(\forall X2.(m2\_subset\_1 X2 k5\_numbers k1\_zf\_lang)\Rightarrow(\forall X3.(\neg v1\_xboole\_0 X3)\Rightarrow((r2\_zf\_model X3 (k11\_zf\_lang (k13\_zf\_lang X2 X0) X1))\Rightarrow(r2\_zf\_model X3 (k11\_zf\_lang X0 X1))))))$$