

t136\_xcplx\_1 (TMKSc-  
CWkuEKZ4kMLr5cBfhAAJnkkJtm5LwY)

October 27, 2020

Let  $v1\_xcplx\_0 : \iota \Rightarrow o$  be given. Let  $k2\_xcplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_xcplx\_0 : \iota \Rightarrow \iota$  be given. Let  $k6\_xcplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xcplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcplx\_0 X1) \Rightarrow (X0 = k6\_xcplx\_0 (k2\_xcplx\_0 X0 X1) X1)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcplx\_0 X0) \wedge (v1\_xcplx\_0 X1)) \Rightarrow (v1\_xcplx\_0 (k2\_xcplx\_0 X0 X1)) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xcplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcplx\_0 X1) \Rightarrow (k6\_xcplx\_0 X0 X1 = k2\_xcplx\_0 X0 (k4\_xcplx\_0 X1))) \quad (3)$$

**Theorem 1**

$$\forall X0.(v1\_xcplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcplx\_0 X1) \Rightarrow (X0 = k2\_xcplx\_0 (k2\_xcplx\_0 X0 X1) (k4\_xcplx\_0 X1)))$$