

t136_xxreal_1 (TMcvEPZvxgMVed- dQLC8PTSJJWPnQooskDwR)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_xxreal_0 X1) \Rightarrow (\forall X2. (v1_xxreal_0 X2) \Rightarrow ((X0 \in k4_xxreal_1 X1 X2) \Rightarrow ((X0 \in k2_xxreal_1 X1 X2) \wedge (X0 \neq X1)))) \quad (1)$$

Assume the following.

$$\forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (k2_xxreal_1 X0 X1 = k2_xboole_0 (k1_tarski X0) (k4_xxreal_1 X0 X1)))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (\neg X0 \in X1) \Rightarrow (k4_xboole_0 (k2_xboole_0 X1 (k1_tarski X0)) (k1_tarski X0) = X1) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k6_subset_1 X0 X1 = k4_xboole_0 X0 X1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k2_xboole_0 X0 X1 = k2_xboole_0 X1 X0 \quad (5)$$

Theorem 1

$$\forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (k6_subset_1 (k2_xxreal_1 X0 X1) (k1_tarski X0) = k4_xxreal_1 X0 X1)))$$