

t138_rvsum_1

(TMag855gDCtM3hHUUjeEmqKujvjYScEncM8)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k4_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k23_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k11_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k4_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v3_card_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 \\ & X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 \\ & X1) \wedge ((v3_valued_0 X1) \wedge (v1_finseq_1 X1)))) \Rightarrow ((k3_finseq_1 X0 = \\ & k3_finseq_1 X1) \Rightarrow (k23_rvsum_1 (k4_rvsum_1 X0 X1) (k4_rvsum_1 X0 \\ & X1) = k9_binop_2 (k9_binop_2 (k23_rvsum_1 X0 X0) (k11_binop_2 np_2 \\ & (k23_rvsum_1 X0 X1))) (k23_rvsum_1 X1 X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_2 X1 X0) \Rightarrow (\forall X2. (m2_finseq_2 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. ((v7_ordinal1 X0) \wedge ((m1_subset_1 \\ & X1 (k4_finseq_2 X0 k1_numbers)) \wedge (m1_subset_1 X2 (k4_finseq_2 \\ & X0 k1_numbers)))) \Rightarrow (k5_rvsum_1 X0 X1 X2 = k1_valued_1 X1 X2) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge((v3_valued_0 X0)\wedge(v1_finseq_1 X0))))\wedge((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge((v3_valued_0 X1)\wedge(v1_finseq_1 X1))))))\Rightarrow(k4_rvsum_1 X0 X1 = k1_valued_1 X0 X1) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_finseq_1 X0)))\Rightarrow(k3_finseq_1 X0 = k1_card_1 X0) \quad (6)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2 X1 X0)\Rightarrow(\forall X2.(m2_finseq_2 X2 X0 X1)\Rightarrow(m2_finseq_1 X2 X0)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow((v1_relat_1 X1)\wedge((v1_funct_1 X1)\wedge(v1_finseq_1 X1))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(v7_ordinal1 X0)\Rightarrow(m1_finseq_2 (k4_finseq_2 X0 X1) X1) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.(v3_card_1 X1 X0)\Leftrightarrow(k1_card_1 X1 = X0) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (13)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v5_relat_1 X0 k1_numbers))\Rightarrow((v1_relat_1 X0)\wedge(v3_valued_0 X0)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge(v7_ordinal1 X1))\Rightarrow(\forall X2.(m1_subset_1 X2 (k4_finseq_2 X1 X0))\Rightarrow(v3_card_1 X2 X1)) \quad (15)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v7_ordinal1\ X0) \Rightarrow (\forall X1.(m2_finseq_2\ X1\ k1_numbers \\ & (k4_finseq_2\ X0\ k1_numbers)) \Rightarrow (\forall X2.(m2_finseq_2\ X2\ k1_numbers \\ & (k4_finseq_2\ X0\ k1_numbers)) \Rightarrow (k23_rvsum_1\ (k5_rvsum_1\ X0\ X1\ X2) \\ & (k5_rvsum_1\ X0\ X1\ X2) = k9_binop_2\ (k9_binop_2\ (k23_rvsum_1\ X1\ X1) \\ & (k11_binop_2\ np_2\ (k23_rvsum_1\ X1\ X2)))\ (k23_rvsum_1\ X2\ X2)))) \end{aligned}$$