

t138_seq_4 (TMQrbAmVCML- TyNNC26XbFKo3CmBiqteiwdY)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v5_valued_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_seq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1)) \quad (1)$$

Assume the following.

$$\forall X0.(m2_finseq_1 X0 k1_numbers) \Rightarrow ((v5_valued_0 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow (((X1 \in k4_finseq_1 X0) \wedge ((X2 \in k4_finseq_1 X0) \wedge (r1_xxreal_0 X1 X2))) \Rightarrow (r1_xxreal_0 (k1_seq_1 X0 X1) (k1_seq_1 X0 X2)))))) \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (4)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Rightarrow(m1_subset_1 X2 X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0)\Rightarrow((v1_funct_1 X1)\wedge((v1_finseq_1 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (7)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (8)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v5_relat_1 X0 k1_numbers)\wedge(v1_funct_1 X0))))\Rightarrow((v5_valued_0 X0)\Leftrightarrow(\forall X1.(m2_subset_1 X1 k1_numbers k5_numbers)\Rightarrow(\forall X2.(m2_subset_1 X2 k1_numbers k5_numbers)\Rightarrow(\neg(X1 \in k9_xtuple_0 X0)\wedge((X2 \in k9_xtuple_0 X0)\wedge((\neg r1_xxreal_0 X2 X1)\wedge(r1_xxreal_0 (k1_seq_1 X0 X2) (k1_seq_1 X0 X1)))))))))) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0)\wedge(v1_xxreal_0 X1))\Rightarrow((r1_xxreal_0 X0 X1)\vee(r1_xxreal_0 X1 X0)) \quad (10)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (13)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0))\Rightarrow(v1_xboole_0 X1)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (15)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 k1_numbers) \Rightarrow ((v5_valued_0 X0) \Rightarrow (\forall X1. \\ & (m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 k1_numbers k5_numbers) \Rightarrow (\neg(X1 \in k4_finseq_1 X0) \wedge ((X2 \in k4_finseq_1 \\ & X0) \wedge ((X1 \neq X2) \wedge (k1_seq_1 X0 X1 = k1_seq_1 X0 X2))))))) \end{aligned}$$