

t139_funct_7 (TMNMNrEeqPX- eAmDJq7VjM4eoJrzcGgXdnjD)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funct_7 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. \forall X2. \\ k9_xtuple_0 (k2_funct_7 X0 X2 X1) = k9_xtuple_0 X0) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1. ((\\ v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge \\ (v1_funct_1 X2)) \Rightarrow (k1_funct_4 (k1_funct_4 X0 X1) X2 = k1_funct_4 \\ X0 (k1_funct_4 X1 X2)))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow (k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k2_funcop_1 X0 X1)) \wedge (v1_funct_1 (k2_funcop_1 X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.(v1_funct_1 (k7_funcop_1 X0 X1))\wedge((v1_funct_2 (k7_funcop_1 X0 X1) X0 (k1_tarSKI X1))\wedge(m1_subset_1 (k7_funcop_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 X0 (k1_tarSKI X1)))))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0)\wedge(v1_funct_1 X0))\wedge((v1_relat_1 X1)\wedge(v1_funct_1 X1)))\Rightarrow((v1_relat_1 (k1_funct_4 X0 X1))\wedge(v1_funct_1 (k1_funct_4 X0 X1))) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.k16_funcop_1 X0 X1 = k7_funcop_1 (k1_tarSKI X0) X1 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.k4_funct_4 X0 X1 X2 X3 = k1_funct_4 (k16_funcop_1 X0 X2) (k16_funcop_1 X1 X3) \quad (10)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge(v1_funct_1 X0))\Rightarrow(\forall X1.\forall X2.((X1 \in k9_xtuple_0 X0)\Rightarrow(k2_funct_7 X0 X1 X2 = k1_funct_4 X0 (k16_funcop_1 X1 X2)))\wedge((\neg X1 \in k9_xtuple_0 X0)\Rightarrow(k2_funct_7 X0 X1 X2 = X0))) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow((v1_partfun1 X1 X0)\Leftrightarrow(k1_relset_1 X0 X1 = X0)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.k2_funcop_1 X0 X1 = k2_zfmisc_1 X0 (k1_tarSKI X1) \quad (13)$$

Theorem 1

$$\forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.((v1_relat_1 X1)\wedge((v4_relat_1 X1 X0)\wedge((v1_funct_1 X1)\wedge(v1_partfun1 X1 X0))))\Rightarrow(\forall X2.(m1_subset_1 X2 X0)\Rightarrow(\forall X3.(m1_subset_1 X3 X0)\Rightarrow(\forall X4.\forall X5.k1_funct_4 X1 (k4_funct_4 X2 X3 X4 X5) = k2_funct_7 (k2_funct_7 X1 X2 X4) X3 X5))))$$