

t13_ami_2 (TMT-
NevnJ8F7bp65SFavx4RABPXMHHz2F7CJ)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_card_3 : \iota \Rightarrow \iota$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_ami_2 : \iota$ be given. Let $k4_ami_2 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_ami_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $k1_funct_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v4_funct_1 : \iota \Rightarrow o$ be given. Let $v2_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_ami_2 : \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k16_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (k9_xtuple_0 (k2_funcop_1 X0 X1) = X0) \wedge (r1_tarski (k10_xtuple_0 (k2_funcop_1 X0 X1)) (k1_tarski X1)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2. ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((\neg X0 \in k9_xtuple_0 X1) \Rightarrow (k1_funct_1 (k1_funct_4 X2 X1) X0 = k1_funct_1 X2 X0))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. k7_funcop_1 X0 X1 = k2_funcop_1 X0 X1 \quad (3)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (4)$$

Assume the following.

$$\neg v1_finset_1 k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (v4_funct_1 (k4_card_3 X0)) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. (((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \wedge ((v1_relat_1 X1) \wedge (v1_funct_1 X1))) \Rightarrow ((v1_relat_1 (k3_relat_1 X0 X1)) \wedge (v1_funct_1 (k3_relat_1 X0 X1))) \quad (7)$$

Assume the following.

$$(v1_relat_1 (k3_relat_1 k3_ami_2 k4_ami_2)) \wedge (v2_relat_1 (k3_relat_1 k3_ami_2 k4_ami_2)) \quad (8)$$

Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 (k2_funcop_1 X0 X1)) \wedge (v1_funct_1 (k2_funcop_1 X0 X1)) \quad (9)$$

Assume the following.

$$(v1_relat_1 k4_ami_2) \wedge ((v4_relat_1 k4_ami_2 np_2) \wedge ((v1_funct_1 k4_ami_2) \wedge (v1_partfun1 k4_ami_2 np_2))) \quad (10)$$

Assume the following.

$$(v1_funct_1 k3_ami_2) \wedge ((v1_funct_2 k3_ami_2 k1_ami_2 np_2) \wedge (m1_subset_1 k3_ami_2 (k1_zfmisc_1 (k2_zfmisc_1 k1_ami_2 np_2)))) \quad (11)$$

Assume the following.

$$\forall X0. \forall X1. k16_funcop_1 X0 X1 = k7_funcop_1 (k1_tarski X0) X1 \quad (12)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 (k4_card_3 (k3_relat_1 k3_ami_2 k4_ami_2))) \Rightarrow (\forall X1. (v7_ordinal1 X1) \Rightarrow (k6_ami_2 X0 X1 = k1_funct_4 X0 (k16_funcop_1 k5_numbers X1))) \quad (13)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (14)$$

Assume the following.

$$\forall X0. (v4_funct_1 X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge (v1_funct_1 X1))) \quad (15)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_finset_1 X0) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (17)$$

Theorem 1

$$\begin{aligned} \forall X0.(m1_subset_1 X0 (k4_card_3 (k3_relat_1 k3_ami_2 k4_ami_2))) \Rightarrow \\ (\forall X1.(v7_ordinal1 X1) \Rightarrow (\forall X2.(v7_ordinal1 X2) \Rightarrow (\\ k1_funct_1 (k6_ami_2 X0 X1) X2 = k1_funct_1 X0 X2))) \end{aligned}$$