

t13_euclid

(TMcJkFsqdNHrL7gRnCePh6oiZ9XNfZC8Zce)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m2_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k12_euclid : \iota \Rightarrow \iota$ be given. Let $k8_euclid : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_euclid : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $k6_rvsum_1 : \iota \Rightarrow \iota$ be given. Let $m1_finseq_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k45_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k30_valued_1 : \iota \Rightarrow \iota$ be given. Let $k6_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (m2_finseq_2 X1 k1_numbers \\ (k1_euclid X0)) \Rightarrow (\forall X2. (m2_finseq_2 X2 k1_numbers (k1_euclid \\ X0)) \Rightarrow (r1_xxreal_0 (k12_euclid (k7_euclid X0 X1 X2)) (k7_real_1 \\ (k12_euclid X1) (k12_euclid X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v1_finseq_1 \\ X0) \wedge (v3_valued_0 X0)))) \Rightarrow (k12_euclid (k6_rvsum_1 X0) = k12_euclid \\ X0) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. (m1_finseq_2 X1 X0) \Rightarrow (\forall X2. (m2_finseq_2 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. ((v7_ordinal1 X0) \wedge ((m1_subset_1 \\ X1 (k1_euclid X0)) \wedge (m1_subset_1 X2 (k1_euclid X0)))) \Rightarrow (k8_euclid \\ X0 X1 X2 = k45_valued_1 X1 X2) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v7_ordinal1\ X0)\wedge((m1_subset_1\ X1\ (k1_euclid\ X0))\wedge(m1_subset_1\ X2\ (k1_euclid\ X0))))\Rightarrow(k7_euclid\ X0\ X1\ X2 = k1_valued_1\ X1\ X2) \quad (5)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge((v3_valued_0\ X0)\wedge(v1_finseq_1\ X0))))\Rightarrow(k6_rvsum_1\ X0 = k30_valued_1\ X0) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(m1_subset_1\ X1\ (k1_euclid\ X0)))\Rightarrow(k6_euclid\ X0\ X1 = k30_valued_1\ X1) \quad (7)$$

Assume the following.

$$v3_membered\ k1_numbers \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_2\ X1\ X0)\Rightarrow(\forall X2.(m2_finseq_2\ X2\ X0\ X1)\Rightarrow(m2_finseq_1\ X2\ X0)) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1\ X1\ X0)\Rightarrow((v1_funct_1\ X1)\wedge((v1_finseq_1\ X1)\wedge(m1_subset_1\ X1\ (k1_zfmisc_1\ (k2_zfmisc_1\ k5_numbers\ X0)))))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(m1_subset_1\ X1\ (k1_euclid\ X0)))\Rightarrow(m2_finseq_2\ (k6_euclid\ X0\ X1)\ k1_numbers\ (k1_euclid\ X0)) \quad (11)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_valued_0\ X0)))\Rightarrow((v1_relat_1\ (k30_valued_1\ X0))\wedge((v1_funct_1\ (k30_valued_1\ X0))\wedge(v1_valued_0\ (k30_valued_1\ X0)))) \quad (12)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(m1_finseq_2\ (k1_euclid\ X0)\ k1_numbers) \quad (13)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge((v1_finseq_1\ X0)\wedge(v3_valued_0\ X0))))\Rightarrow(m1_subset_1\ (k12_euclid\ X0)\ k1_numbers) \quad (14)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 X0))) \Rightarrow \\ (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_valued_0 \\ X1)))) \Rightarrow (k45_valued_1 X0 X1 = k1_valued_1 X0 (k30_valued_1 X1))) \end{aligned} \quad (15)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 X0 k1_numbers) \wedge (v1_xreal_0 X1)) \Rightarrow (k7_real_1 X0 X1 = k7_real_1 X1 X0) \quad (16)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. (((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_valued_0 \\ X0))) \wedge ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_valued_0 X1)))) \Rightarrow \\ (k1_valued_1 X0 X1 = k1_valued_1 X1 X0) \end{aligned} \quad (17)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \Rightarrow ((v1_relat_1 X0) \wedge (v1_valued_0 X0)) \quad (18)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v5_relat_1 X0 k1_numbers)) \Rightarrow ((v1_relat_1 X0) \wedge (v3_valued_0 X0)) \quad (19)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (20)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (21)$$

Assume the following.

$$\forall X0. (v3_membered X0) \Rightarrow (\forall X1. (m1_subset_1 X1 X0) \Rightarrow (v1_xreal_0 X1)) \quad (22)$$

Theorem 1

$$\begin{aligned} \forall X0. (v7_ordinal1 X0) \Rightarrow (\forall X1. (m2_finseq_2 X1 k1_numbers \\ (k1_euclid X0)) \Rightarrow (\forall X2. (m2_finseq_2 X2 k1_numbers (k1_euclid \\ X0)) \Rightarrow (r1_xxreal_0 (k12_euclid (k8_euclid X0 X1 X2)) (k7_real_1 \\ (k12_euclid X1) (k12_euclid X2)))))) \end{aligned}$$