

t13_frechet

(TMZhqzjVzH6fmdd8NaiuAeUCR1bDuYbWJCn)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_partfun1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_relat_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (k7_relat_1 (k4_relat_1 X0) X1 = X1) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 (k1_zfmisc_1 X1)) \Leftrightarrow (r1_tarski X0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (k7_relset_1 X0 X1 X2 X3 = k7_relat_1 X2 X3) \quad (3)$$

Assume the following.

$$\forall X0. k6_partfun1 X0 = k4_relat_1 X0 \quad (4)$$

Assume the following.

$$\forall X0. (v1_partfun1 (k6_partfun1 X0) X0) \wedge (m1_subset_1 (k6_partfun1 X0) (k1_zfmisc_1 (k2_zfmisc_1 X0 X0))) \quad (5)$$

Theorem 1

$$\forall X0. \forall X1. (r1_tarski X1 X0) \Rightarrow (k7_relset_1 X0 X0 (k6_partfun1 X0) X1 = X1)$$