

t13\_group\_7 (TMN-  
tjr6QAL51suG8ZgZmXPpHNZeta8JFBmD)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v5\_group\_1 : \iota \Rightarrow o$  be given. Let  $l3\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k9\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $np\_1 : \iota$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_group\_7 : \iota \Rightarrow o$  be given. Let  $v4\_group\_7 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k5\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_group\_7 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_tarski : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k2\_finseq\_1 : \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0. \forall X1. (m1\_subset\_1 X0 X1) \Rightarrow ((v1\_xboole\_0 X1) \vee (X0 \in X1)) \quad (1)$$

Assume the following.

$$\forall X0. ((\neg v2\_struct\_0 X0) \wedge (l3\_algstr\_0 X0)) \Rightarrow ((v1\_relat\_1 (k9\_finseq\_1 X0)) \wedge ((v4\_relat\_1 (k9\_finseq\_1 X0) (k1\_tarski np\_1)) \wedge ((v1\_funct\_1 (k9\_finseq\_1 X0)) \wedge ((v1\_partfun1 (k9\_finseq\_1 X0) (k1\_tarski np\_1)) \wedge (v1\_group\_7 (k9\_finseq\_1 X0)))))) \quad (2)$$

Assume the following.

$$\forall X0. k9\_finseq\_1 X0 = k5\_finseq\_1 X0 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. ((\neg v1\_xboole\_0 X0) \wedge (((v1\_relat\_1 X1) \wedge ((v4\_relat\_1 X1 X0) \wedge ((v1\_funct\_1 X1) \wedge ((v1\_partfun1 X1 X0) \wedge (v1\_group\_7 X1)))))) \wedge (m1\_subset\_1 X2 X0))) \Rightarrow (k1\_group\_7 X0 X1 X2 = k1\_funct\_1 X1 X2) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. v1\_relat\_1 (k1\_tarski (k4\_tarski X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0. \neg v1\_xboole\_0 (k1\_tarski X0) \quad (6)$$

Assume the following.

$$\forall X0. \forall X1. v1\_funct\_1 (k1\_tarski (k4\_tarski X0 X1)) \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. ((v1\_relat\_1 X1) \wedge ( \\ (v4\_relat\_1 X1 X0) \wedge ((v1\_funct\_1 X1) \wedge ((v1\_partfun1 X1 X0) \wedge (v1\_group\_7 \\ X1)))))) \Rightarrow ((v4\_group\_7 X1 X0) \Leftrightarrow (\forall X2. (m1\_subset\_1 X2 X0) \Rightarrow \\ (v5\_group\_1 (k1\_group\_7 X0 X1 X2)))))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((X1 = \\ k9\_finseq\_1 X0) \Leftrightarrow ((k9\_xtuple\_0 X1 = k2\_finseq\_1 np\_1) \wedge (k1\_funct\_1 \\ X1 np\_1 = X0))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0. k5\_finseq\_1 X0 = k1\_tarski (k4\_tarski np\_1 X0) \quad (10)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1\_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (11)$$

### Theorem 1

$$\begin{aligned} \forall X0. ((\neg v2\_struct\_0 X0) \wedge ((v5\_group\_1 X0) \wedge (l3\_algstr\_0 \\ X0))) \Rightarrow ((v1\_relat\_1 (k9\_finseq\_1 X0)) \wedge ((v4\_relat\_1 (k9\_finseq\_1 \\ X0) (k1\_tarski np\_1)) \wedge ((v1\_funct\_1 (k9\_finseq\_1 X0)) \wedge ((v1\_partfun1 \\ (k9\_finseq\_1 X0) (k1\_tarski np\_1)) \wedge ((v1\_group\_7 (k9\_finseq\_1 \\ X0)) \wedge (v4\_group\_7 (k9\_finseq\_1 X0) (k1\_tarski np\_1))))))) \end{aligned}$$