

t13_matrixr1

(TMZyyD3XVU7hn6Cq77xf21NRWPRhJsRphyM)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k8_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v3_valued_0 \\ & X0) \wedge (v1_finseq_1 X0)))) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 \\ & X1) \wedge ((v3_valued_0 X1) \wedge (v1_finseq_1 X1)))) \Rightarrow (\forall X2.((v1_relat_1 \\ & X2) \wedge ((v1_funct_1 X2) \wedge ((v3_valued_0 X2) \wedge (v1_finseq_1 X2)))) \Rightarrow \\ & (k8_rvsum_1 X0 (k8_rvsum_1 X1 X2) = k4_rvsum_1 (k8_rvsum_1 X0 X1 \\ & X2))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge \\ & (v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers \\ & X0)))) \end{aligned} \tag{3}$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge (v1_funct_1 X1) \wedge (v1_finseq_1 X1)) \tag{4}$$

Assume the following.

$$\forall X0. (m1_finseq_1 X0 k1_numbers) \Rightarrow (v3_valued_0 X0) \tag{5}$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 k1_numbers) \Rightarrow (\forall X1.(m2_finseq_1 \\ & X1 k1_numbers) \Rightarrow (\forall X2.(m2_finseq_1 X2 k1_numbers) \Rightarrow (((k3_finseq_1 \\ & X0 = k3_finseq_1 X1) \wedge (k3_finseq_1 X1 = k3_finseq_1 X2)) \Rightarrow (k8_rsum_1 \\ & X0 (k8_rsum_1 X1 X2) = k4_rsum_1 (k8_rsum_1 X0 X1) X2)))) \end{aligned}$$