

t13_metric_2 (TM- RMpkj8nwNzTT6T2S9X4xSYT25rhd2gLY)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v6_metric_1 : \iota \Rightarrow o$ be given. Let $v7_metric_1 : \iota \Rightarrow o$ be given. Let $v8_metric_1 : \iota \Rightarrow o$ be given. Let $v9_metric_1 : \iota \Rightarrow o$ be given. Let $l1_metric_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_metric_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $r1_metric_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v6_metric_1 X0) \wedge ((v7_metric_1 \\ & X0) \wedge ((v8_metric_1 X0) \wedge ((v9_metric_1 X0) \wedge (l1_metric_1 X0)))))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 \\ & X2 (u1_struct_0 X0)) \Rightarrow ((X2 \in k1_metric_2 X0 X1) \Leftrightarrow (X2 = X1)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k6_domain_1 X0 X1 = k1_tarski X1) \quad (2)$$

Assume the following.

$$\forall X0. ((\neg v2_struct_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\neg v1_xboole_0 (u1_struct_0 X0)) \quad (3)$$

Assume the following.

$$\forall X0. (l1_metric_1 X0) \Rightarrow (l1_struct_0 X0) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge (l1_metric_1 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (k1_metric_2 X0 X1 = \text{ReplSep} \\ & (\text{toSet } (\lambda X2 : \iota. m1_subset_1 X2 (u1_struct_0 X0))) (\lambda X2 : \\ & \iota. r1_metric_2 X0 X1 X2) (\lambda X2 : \iota. X2))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k1_tarski X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (X2 = X0)) \quad (6)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v6_metric_1 X0) \wedge ((v7_metric_1 \\ & X0) \wedge ((v8_metric_1 X0) \wedge ((v9_metric_1 X0) \wedge (l1_metric_1 X0)))))) \Rightarrow \\ & (\forall X1. (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow (k1_metric_2 X0 \\ & X1 = k6_domain_1 (u1_struct_0 X0) X1)) \end{aligned}$$