

# t13\_nagata\_2 (TMQtisNr- WELMX6SQwGC9MWJfCVZHrauZWei)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k5\_numbers : \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k18\_complex1 : \iota \Rightarrow \iota$  be given. Let  $k8\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_series\_1 : \iota \Rightarrow \iota$  be given. Let  $k56\_valued\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_seq\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v1\_membered : \iota \Rightarrow o$  be given. Let  $k54\_valued\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v3\_valued\_0 : \iota \Rightarrow o$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v3\_membered : \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0.((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers k1\_numbers) \wedge \\ & (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))))) \Rightarrow \\ & (\forall X1.(m1\_subset\_1 X1 k5\_numbers) \Rightarrow (r1\_xxreal\_0 (k18\_complex1 \\ & (k1\_seq\_1 (k3\_series\_1 X0) X1)) (k1\_seq\_1 (k3\_series\_1 (k56\_valued\_1 \\ & k5\_numbers k1\_numbers X0)) X1))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\ & (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\ & X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1\_funct\_1 X1) \wedge ((v1\_funct\_2 \\ & X1 k5\_numbers X0) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers \\ & X0)))))) \wedge (v7\_ordinal1 X2)) \Rightarrow (k8\_nat\_1 X0 X1 X2 = k1\_funct\_1 X1 X2) \end{aligned} \tag{3}$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1\_membered\ X1)\wedge((v1\_funct\_1\ X2)\wedge(m1\_subset\_1\ X2\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ X0\ X1)))))\Rightarrow(k56\_valued\_1\ X0\ X1\ X2 = k54\_valued\_1\ X2)$$
(5)

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1\ X0)\wedge((v1\_funct\_1\ X0)\wedge(v3\_valued\_0\ X0)))\Rightarrow(k1\_seq\_1\ X0\ X1 = k1\_funct\_1\ X0\ X1)$$
(6)

Assume the following.

$$\forall X0.\forall X1.\forall X2.(((\neg v1\_xboole\_0\ X1)\wedge(v1\_membered\ X1))\wedge((v1\_funct\_1\ X2)\wedge((v1\_funct\_2\ X2\ X0\ X1)\wedge(m1\_subset\_1\ X2\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ X0\ X1))))))\Rightarrow((v1\_funct\_1\ (k54\_valued\_1\ X2))\wedge(v1\_partfun1\ (k54\_valued\_1\ X2)\ X0))$$
(7)

Assume the following.

$$(\neg v1\_xboole\_0\ k4\_ordinal1)\wedge(v3\_ordinal1\ k4\_ordinal1)$$
(8)

Assume the following.

$$v3\_membered\ k1\_numbers$$
(9)

Assume the following.

$$\neg v1\_xboole\_0\ k1\_numbers$$
(10)

Assume the following.

$$m1\_subset\_1\ k5\_numbers\ (k1\_zfmisc\_1\ k1\_numbers)$$
(11)

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1\_membered\ X1)\wedge((v1\_funct\_1\ X2)\wedge(m1\_subset\_1\ X2\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ X0\ X1)))))\Rightarrow((v1\_funct\_1\ (k56\_valued\_1\ X0\ X1\ X2))\wedge(m1\_subset\_1\ (k56\_valued\_1\ X0\ X1\ X2)\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ X0\ k1\_numbers))))$$
(12)

Assume the following.

$$\forall X0.((v1\_funct\_1\ X0)\wedge((v1\_funct\_2\ X0\ k5\_numbers\ k1\_numbers)\wedge(m1\_subset\_1\ X0\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ k5\_numbers\ k1\_numbers)))))\Rightarrow((v1\_funct\_1\ (k3\_series\_1\ X0))\wedge((v1\_funct\_2\ (k3\_series\_1\ X0\ k5\_numbers\ k1\_numbers)\wedge(m1\_subset\_1\ (k3\_series\_1\ X0)\ (k1\_zfmisc\_1\ (k2\_zfmisc\_1\ k5\_numbers\ k1\_numbers))))))$$
(13)

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (14)$$

Assume the following.

$$\forall X0.(v3\_membered X0) \Rightarrow (v1\_membered X0) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \quad (16)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v1\_partfun1 X2 X0) \Rightarrow (v1\_funct\_2 X2 X0 X1)) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.(v3\_membered X1) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v3\_valued\_0 X2)) \quad (18)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((v1\_funct\_1 X0) \wedge ((v1\_funct\_2 X0 k5\_numbers k1\_numbers) \wedge \\ & (m1\_subset\_1 X0 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k5\_numbers k1\_numbers)))))) \Rightarrow \\ & (\forall X1.(m2\_subset\_1 X1 k1\_numbers k5\_numbers) \Rightarrow (r1\_xxreal\_0 \\ & (k18\_complex1 (k8\_nat\_1 k1\_numbers (k3\_series\_1 X0) X1) (k8\_nat\_1 \\ & k1\_numbers (k3\_series\_1 (k56\_valued\_1 k5\_numbers k1\_numbers \\ & X0)) X1))) \end{aligned}$$