

t13_orders_1

(TMR61zDz4a2zbx2K3UmvB9MvXTKh3dKJ8Gc)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_relat_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_relat_1 : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 X0))) \Rightarrow ((\forall X3.\neg(X3 \in X1) \wedge (\forall X4.\neg k4_tarski X3 X4 \in X2)) \Leftrightarrow (k1_relset_1 X1 X2 = X1)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v1_partfun1 X1 X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 X0)))) \Rightarrow (k1_relat_1 X1 = X0) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.k4_tarski X0 X1 = k2_tarski (k2_tarski X0 X1) (k1_tarski X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v4_relat_1 X1 X0)) \Rightarrow ((v1_partfun1 X1 X0) \Leftrightarrow (k1_relset_1 X0 X1 = X0)) \quad (4)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.(r1_relat_2 X0 X1) \Leftrightarrow (\forall X2.(X2 \in X1) \Rightarrow (k4_tarski X2 X2 \in X0))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (7)$$

Theorem 1

$$\forall X0.\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 X0 X0)))\Rightarrow((r1_relat_2 X1 X0)\Rightarrow((k1_relset_1 X0 X1 = X0)\wedge(k1_relat_1 X1 = X0)))$$