

t13_rat_1 (TMYAGh-
wjHgmP2YDoUtAwL4GRQWAZT8gNYE3)

October 27, 2020

Let $v1_rat_1 : \iota \Rightarrow o$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_real_1 : \iota \Rightarrow \iota$ be given. Let $k1_rat_1 : \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k5_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xreal_0 : \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow ((r1_xreal_0 np_1 X0) \Rightarrow (r1_xreal_0 (k5_xcmplx_0 X0) np_1)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_rat_1 X0) \Rightarrow (r1_xreal_0 np_1 (k1_rat_1 X0)) \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (k2_real_1 X0 = k5_xcmplx_0 X0) \quad (4)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 X2 X0 X1) \Rightarrow (m1_subset_1 X2 X0)) \quad (6)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (7)$$

Assume the following.

$$\forall X0.(v1_rat_1 X0) \Rightarrow (m2_subset_1 (k1_rat_1 X0) k1_numbers \quad k5_numbers) \quad (8)$$

Assume the following.

$$k1_xxreal_0 = k1_numbers \quad (9)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (10)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (11)$$

Theorem 1

$$\forall X0.(v1_rat_1 X0) \Rightarrow (r1_xxreal_0 (k2_real_1 (k1_rat_1 X0)) \quad np_1)$$