

t13\_rlvect\_x  
(TMF4s1WXUFgmp5jZpXNDT5FB44Umuq8dM1L)

October 27, 2020

Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $l1\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k2\_rlvect\_x : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $m2\_rlvect\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k6\_rlvect\_2 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k4\_numbers : \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0. (&(\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\ &X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\ &((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge (l1\_rlvect\_1 \\ &X0)))))))))) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 \\ &X0))) \Rightarrow (\forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (u1\_struct\_0 \\ &X0))) \Rightarrow (\forall X3. (m2\_rlvect\_2 X3 X0 X1) \Rightarrow ((r1\_tarski X1 X2) \Rightarrow ( \\ &m2\_rlvect\_2 X3 X0 X2)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1\_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0. (&(\neg v2\_struct\_0 X0) \wedge ((v13\_algstr\_0 X0) \wedge ((v2\_rlvect\_1 \\ &X0) \wedge ((v3\_rlvect\_1 X0) \wedge ((v4\_rlvect\_1 X0) \wedge ((v5\_rlvect\_1 X0) \wedge \\ &((v6\_rlvect\_1 X0) \wedge ((v7\_rlvect\_1 X0) \wedge ((v8\_rlvect\_1 X0) \wedge (l1\_rlvect\_1 \\ &X0)))))))))) \Rightarrow (\forall X1. (m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 \\ &X0))) \Rightarrow (k2\_rlvect\_x X0 X1 = ReplSep (toset (\lambda X2 : \iota. m2\_rlvect\_2 \\ &X2 X0 X1)) (\lambda X2 : \iota. r1\_tarski (k2\_relset\_1 k1\_numbers X2) k4\_numbers) \\ &(\lambda X2 : \iota. k6\_rlvect\_2 X0 X2))) \end{aligned} \quad (3)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.((\neg v2\_struct\_0 X0) \wedge (v13\_algstr\_0 X0) \wedge (v2\_rlvect\_1 \\ & X0) \wedge (v3\_rlvect\_1 X0) \wedge (v4\_rlvect\_1 X0) \wedge (v5\_rlvect\_1 X0) \wedge \\ & ((v6\_rlvect\_1 X0) \wedge (v7\_rlvect\_1 X0) \wedge (v8\_rlvect\_1 X0) \wedge (l1\_rlvect\_1 \\ & X0)))))) \Rightarrow (\forall X1.(m1\_subset\_1 X1 (k1\_zfmisc\_1 (u1\_struct\_0 \\ & X0))) \Rightarrow (\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (u1\_struct\_0 \\ & X0))) \Rightarrow ((r1\_tarski X1 X2) \Rightarrow (r1\_tarski (k2\_rlvect\_x X0 X1) (k2\_rlvect\_x \\ & X0 X2)))))) \end{aligned}$$